



# Managing **PSORIATIC ARTHRITIS** in Specialty Practice: *New Therapies, Guidelines and Treatment Targets During the COVID-19 Pandemic*



**THURSDAY  
JANUARY 21, 2021**

**FACULTY**

**Daniel Furst, MD**

Professor of Rheumatology and Medicine  
University of California, Los Angeles, CA  
University of Washington, Seattle, WA  
University of Florence, Florence, Italy



*This activity is provided by Med Learning Group.  
This activity is co-provided by Ultimate Medical Academy/Complete Conference Management (CCM).*

*This activity is supported by an educational grant from Lilly.*

***The RELIEF Initiative***  
***Managing Psoriatic Arthritis in Specialty Practice:***  
***New Therapies, Guidelines and Treatment Targets During the COVID-19 Pandemic***

**FACULTY**

**Daniel E. Furst, MD**

Professor of Rheumatology and Medicine  
University of California, Los Angeles, CA  
University of Washington, Seattle, WA  
University of Florence, Florence, Italy

**PROGRAM OVERVIEW**

This case-based live virtual activity will cover the treatment and management of patients with psoriatic arthritis.

**TARGET AUDIENCE**

This activity is intended for rheumatologists and rheumatology advanced practice providers (NPs and PAs) who are involved in the care and treatment of patients with psoriatic arthritis.

**Learning Objectives**

- Identify the risk of COVID-19-related infections in PsA, along with their impact on therapeutic choice
- Pursue strategies to optimize PsA therapy in the COVID-19 era while minimizing risks and adverse events
- Assess methods for better evaluating and communicating with patients through telemedicine and virtual platforms
- Apply new ways to initiate and manage PsA treatment, monitor PsA disease progression and address adverse events via virtual communication

**ACCREDITATION STATEMENT**

Med Learning Group is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

This CME activity was planned and produced in accordance with the ACCME Essentials.

**CREDIT DESIGNATION STATEMENT**

Med Learning Group designates this live virtual activity for a maximum of 1.0 *AMA Category 1 Credit*<sup>™</sup>. Physicians should claim only the credit commensurate with the extent of their participation in the live virtual activity.

**NURSING CREDIT INFORMATION**

Purpose: This program would be beneficial for nurses involved in the care of patients with psoriatic arthritis.

**CNE Credits:** 1.0 ANCC Contact Hour.

**CNE ACCREDITATION STATEMENT**

Ultimate Medical Academy/CCM is accredited as a provider of continuing nursing education by the American Nurses Credentialing Center's Commission on Accreditation. Awarded 1.0 contact hour of continuing nursing education of RNs and APNs.

## **ABIM Maintenance of Certification**

Successful completion of this CME activity, which includes participation in the evaluation component, enables the participant to earn up to 1.0 Medical Knowledge MOC point in the American Board of Internal Medicine's (ABIM) Maintenance of Certification (MOC) program. It is the CME activity provider's responsibility to submit participant completion information to ACCME for the purpose of granting ABIM MOC credit.

## **DISCLOSURE POLICY STATEMENT**

In accordance with the Accreditation Council for Continuing Medical Education (ACCME) Standards for Commercial Support, educational programs sponsored by Med Learning Group must demonstrate balance, independence, objectivity, and scientific rigor. All faculty, authors, editors, staff, and planning committee members participating in an MLG-sponsored activity are required to disclose any relevant financial interest or other relationship with the manufacturers of any commercial products and/or providers of commercial services that are discussed in an educational activity.

## **DISCLOSURE OF CONFLICTS OF INTEREST**

**Daniel E. Furst, MD**, has received grant/research support from Actelion, Amgen, BMS, Corbus, Galapagos GSK, NIH, Novartis, Pfizer, and Sanofi. He is a consultant with Actelion, Amgen, BMS, Corbus, Galapagos Novartis, and Pfizer.

## **CME Content Review**

The content of this activity was independently peer-reviewed.

The reviewer of this activity has nothing to disclose.

## **CNE Content Review**

The content of this activity was peer-reviewed by a nurse reviewer.

The reviewer of this activity has nothing to disclose.

The staff, planners and managers reported the following financial relationships or relationships to products or devices they or their spouse/life partner have with commercial interests related to the content of this CME/CE activity:

- Matthew Frese, General Manager of Med Learning Group, has nothing to disclose.
- Christina Gallo, SVP, Educational Development of Med Learning Group, has nothing to disclose.
- Ana Maria Albino, Senior Program Manager of Med Learning Group, has nothing to disclose.
- Diana Tommasi, Medical Director of Med Learning Group, has nothing to disclose.
- Lauren Welch, MA, VP of Accreditation and Outcomes of Med Learning Group, has nothing to disclose.
- Brianna Hanson, Accreditation and Outcomes Coordinator of Med Learning Group, has nothing to disclose.

## **DISCLOSURE OF UNLABELED USE**

Med Learning Group requires that faculty participating in any CME activity disclose to the audience when discussing any unlabeled or investigational use of any commercial product or device not yet approved for use in the United States.

During this lecture, faculty may mention the use of medications for both FDA-approved and non-approved indications.

## METHOD OF PARTICIPATION

There are no fees for participating and receiving CME/CE credit for this live virtual activity. To receive CME/CE credit participants must:

1. Read the CME/CNE information and faculty disclosures.
2. Participate in the live virtual activity.
3. Submit the evaluation form to Med Learning Group.

You will receive your certificate upon completion as a downloadable file.

## DISCLAIMER

Med Learning Group makes every effort to develop CME activities that are science-based. This activity is designed for educational purposes. Participants have a responsibility to use this information to enhance their professional development in an effort to improve patient outcomes. Conclusions drawn by the participants should be derived from careful consideration of all available scientific information. The participant should use his/her clinical judgment, knowledge, experience, and diagnostic decision-making before applying any information, whether provided here or by others, for any professional use.

For CME questions, please contact Med Learning Group at [info@medlearninggroup.com](mailto:info@medlearninggroup.com)  
Contact this CME provider at Med Learning Group for privacy and confidentiality policy statement information at [www.medlearninggroup.com/privacy-policy/](http://www.medlearninggroup.com/privacy-policy/)

## AMERICANS WITH DISABILITIES ACT

Staff will be glad to assist you with any special needs. Please contact Med Learning Group prior to participating at [info@medlearninggroup.com](mailto:info@medlearninggroup.com)



This activity is provided by Med Learning Group.



This activity is co-provided by Ultimate Medical Academy/Complete Conference Management (CCM).

Supported by an educational grant from Lilly.



## AGENDA

### Introduction/Background

- Epidemiology and pathophysiology
- COVID-19 background
- COVID-19 and rheumatology: implications for assessment and evaluation
- Disease domains and joint manifestations
- Identification and diagnosis

### Medical Management of PsA in the COVID-19 Era

- 2019 ACR guidelines and their application to practice
- Therapeutic considerations in COVID-19
- Conventional agents
- Biologics, small molecules:
  - Inhibitors of TNF, IL-12/23, IL-17A, IL-23, phosphodiesterase 4, T cell costimulation, and janus kinases
- Evolving standards of treatment in the COVID-19 era
- Treating-to-target: establishing goals of therapy

### Telemedicine and Patient Considerations in the COVID-19 Pandemic

- Early diagnosis and initiation of treatment for long-term success
- Effect of management decisions on patient QoL
- Lowering disease burden (personal, societal, economic) through effective treatment
- Telemedicine: the changing face of rheumatology consults during the pandemic
- Case Study(s)

### Conclusions and Q/A

# *Managing Psoriatic Arthritis in Specialty Practice: New Therapies, Guidelines and Treatment Targets During the COVID-19 Pandemic*

**Daniel E. Furst, MD**

Professor of Rheumatology  
University of California in Los Angeles  
University of Washington, Seattle, Washington  
University of Florence, Florence, Italy

2

## **Disclosures**

- Dr. Furst reports grant/research support from Pfizer, CorbusCSL, Behring, Galapagos, and Mitsubishi. He is a consultant for AbbVie, Amgen, Novartis, Pfizer, and Sanofi.
- During the course of this lecture, Dr Furst will discuss the use of medications for both FDA-approved and non-approved indications

**This educational activity was supported by an educational grant from Lilly.**

3


## Learning Objectives

- Identify the risk of COVID-19-related infections in psoriatic arthritis (PsA), along with their impact on therapeutic choice
- Pursue strategies to optimize PsA therapy in the COVID-19 era while minimizing risks and adverse events
- Assess methods for better evaluating and communicating with patients through telemedicine and virtual platforms
- Apply new ways to initiate and manage PsA treatment, monitor PsA disease progression and address adverse events via virtual communication

4

## COVID-19-associated Hyperinflammation

- Clinical deterioration in COVID-19 often occurs 7-10 days after symptom onset when viral titres decline
  - Pathology likely driven by inflammation rather than direct viral injury
- Elevated inflammatory markers in COVID-19 patients are significantly associated with risk of next-day escalation of respiratory support or death (HR, 2.24)



Manson JJ, et al. *Lancet Rheumatol.* 2020;2:e594-e602. Yang L, et al. *Signal Transduct Target Ther.* 2020;5:128.

5

## Concerns During the COVID-19 Pandemic

- Patients with PsA are not at increased risk of death, invasive ventilation, ICU admission, or serious complications from COVID-19
  - Impact of PsA therapies on COVID-19 disease severity is unknown
- Risk of poor outcomes from COVID-19 appears to be related primarily to general risk factors such as age and comorbidities

Characteristics	Deaths (%)
All confirmed cases	2.3
• Critical cases	49.0
• ≥80 years of age	14.8
• Cardiovascular disease	10.5
• 70-79 years of age	8.0
• Diabetes	9.2
• Chronic respiratory disease	8.0
• Hypertension	6.0
• Cancer	7.6

Mikuls TR, et al. *Arthritis Rheumatol.* 2020;72:1241-1251. Pablos JL, et al. *Ann Rheum Dis.* 2020;79:1544-1549. Wu Z, et al. *JAMA.* 2020;323:1239-1242. Wollina U, et al. *Dermatol Ther.* 2020;33:e13743.

6

## Impact of PsA Comorbidities on COVID-19 Outcomes

- PsA is associated with a higher incidence of CV disease, metabolic syndrome, obesity, diabetes, dyslipidemia, and IBD
- Older age, male sex, and previous comorbidity increased the risk of severe COVID-19 in patients with rheumatic disease and nonrheumatic disease
  - Diabetes and CV disease were associated with a significantly increased risk of severe COVID-19 in rheumatic patients compared to nonrheumatic patients

Variable	Relative risk (95% CI)		P value
	Non-rheumatic cohort	Rheumatic cohort	
Age over 60 years	3.70 (1.99 to 6.93)	4.04 (2.30 to 7.08)	0.841
Male sex	2.16 (1.39 to 3.35)	1.58 (1.09 to 2.29)	0.286
Obesity	1.22 (0.72 to 2.06)	1.62 (1.10 to 2.36)	0.393
Diabetes	0.95 (0.53 to 1.70)	<b>1.93 (1.34 to 2.79)</b>	0.038
Hypertension	1.64 (1.07 to 2.53)	2.27 (1.49 to 3.46)	0.290
CV disease	1.44 (0.90 to 2.33)	<b>2.92 (2.04 to 4.17)</b>	0.020
Lung disease	1.57 (1.00 to 2.46)	1.74 (1.19 to 2.55)	0.723

CV = cardiovascular; IBD = inflammatory bowel disease  
Pablos JL, et al. *Ann Rheum Dis.* 2020;79:1544-1549.

7



## COVID-19 Treatment Modifications

Treatment of Rheumatic Disease in the Absence of COVID-19 Infection or Exposure	
HCQ/CQ, SSZ, MTX, LEF, immunosuppressants (tacrolimus, CSA, MMF, AZA), biologics, JAK inhibitors, NSAIDs	Continue therapy
Low-dose corticosteroids	May be started if clinically indicated (<10 mg prednisone equivalent/day)
Following SARS-CoV-2 Exposure	
HCQ/CQ, SSZ, NSAIDs	May be continued
Immunosuppressants (tacrolimus, CSA, MMF, AZA), non-IL-6 biologics, JAK inhibitors	Stop therapy temporarily, pending a negative COVID-19 test or 2 weeks of symptom-free observation
IL-6 inhibitors	May be continued in select circumstances
Documented or presumptive COVID-19	
HCQ/CQ	May be continued
SSZ, MTX, LEF, non-IL-6 biologics, immunosuppressants, and JAK inhibitors	Withhold or stop therapy
NSAIDs	Should be stopped in patients with severe respiratory symptoms

AZA = azathioprine; CSA = cyclosporine A; CQ = chloroquine; HCQ = hydroxychloroquine; IL = interleukin; JAK = Janus kinase; LEF = leflunomide; MMF = mycophenolate mofetil; MTX = methotrexate; NSAID = nonsteroidal anti-inflammatory drug; SSZ = sulfasalazine  
Mikuls TR, et al. *Arthritis Rheumatol.* 2020;72:1241-1251.

- All recommendations based on very low quality of evidence and moderate to high consensus
- Recommendations are for rheumatic disease in general and are not subdivided by patient disease. There are no specific recommendations for PsA.
  - May reinstate therapy within 7-14 days of symptom resolution for those with mild COVID-19
  - Consider reinitiating therapy in 10-17 days after positive PCR results if asymptomatic COVID-19
  - Timing of reinitiating therapy after severe COVID-19 should be made on case-by-case basis

8

## Corticosteroid Use During the COVID-19 Pandemic

- A case series of 600 patients found prednisone  $\geq 10$  mg/day was associated with increased odds of hospitalization (OR, 2.05)
- A study in patients with inflammatory bowel disease and COVID-19 found steroids increase the risk of severe COVID-19 (aOR, 6.0)

**Glucocorticoids should be used at the lowest possible dose to control rheumatic disease, regardless of exposure or infection status**


**Glucocorticoids should not be abruptly stopped, regardless of exposure or infection status**

OR = odds ratio; aOR = adjusted odds ratio.

Gianfrancesco M, et al. *Ann Rheum Dis.* 2020;79:859-866. Brenner EJ, et al. *Gastroenterology.* 2020;159:481-491. Mikuls TR, et al. *Arthritis Rheumatol.* 2020;72:1241-1251.

9

## Need for Continuity of Care During the COVID-19 Pandemic

- In a study of 1,517 patients in the US with PsA, RA, SpA, or SLE, 14.9% stopped using their DMARD between March-May 2020
- Of the patients who stopped their DMARDs, what percentage of these interruptions were NOT recommended by a physician?  **78.7%**
- 29.5% of patients used telehealth services
  - Treatment interruption was more common among patients who reported that telehealth was not available (25.4% vs 13.1%, respectively)

DMARD = disease-modifying anti-rheumatic drug; RA = rheumatoid arthritis; SLE = systemic lupus erythematosus; SpA = spondyloarthritis.  
George M, et al. *J Rheumatol*. 2020. doi:10.3899/jrheum.201017.

10

## Incorporating Telehealth into Your Practice

- **Schedule enough time.** Telehealth consults often take longer than expected to find the required information
- **Train staff in triaging symptom burden.** Identify patients with unstable symptoms who require an in-person appointment
- **Educate on self-management.** Patients may not come in for a follow-up appointment for weeks or months.
  - Teach about warning signs that require prompt evaluation
  - Educate about how to manage symptoms remotely
  - Ensure patients have enough medication
- **Clarify expectations** of what can or cannot be done remotely
  - Recognize patients who require in-person evaluation

Centers for Disease Control and Prevention website (<https://www.cdc.gov/coronavirus/2019-ncov/hcp/telehealth.html>). Landewe RBM, et al. *Ann Rheum Dis*. 2020;79:851-858.

11

## Strategies to Increase Telehealth Uptake

- **Use technology** that allows you to send and receive patient-reported outcomes scales
- **Prescreen patients with disease activity scales** and request in-person visit if scores are high
- **Offer flexibility** in platforms that can be used for video consultation, and non-video options to serve patients with limited technology and connectivity
- **Postpone** regular blood monitoring and face-to-face consultations in patients with stable disease and therapy without signs of drug toxicity
- **Communicate with insurers/payers** to understand availability of covered telehealth services

Centers for Disease Control and Prevention website (<https://www.cdc.gov/coronavirus/2019-ncov/hcp/telehealth.html>). Landewe RBM, et al. *Ann Rheum Dis.* 2020;79:851-858.

12

## Treatment Options

13

## Initiating Therapy

**Treatment-naïve Active PsA**

Start TNFi biologic over OSM, IL-17i biologic or IL-12/23i biologic

May consider alternative choices in some situations

Start OSM over IL-17i biologic or IL-12/23i biologic

May consider alternative choices in some situations

Start MTX over NSAIDs

May consider alternative choices in some situations

Start IL-17i biologic over IL-12/23i biologic

May consider alternative choices in some situations

Discuss with the patient, since all recommendations are conditional based on low to very low quality evidence

- Contraindications to TNFi include congestive heart failure, previous serious infection, recurrent infections, or demyelinating disease
- An OSM (MTX, SSZ, LEF, CSA, or APR) may be considered if disease is not severe, oral therapy is preferred, or patient does not want to start a biologic

APR = apremilast; CSA = cyclosporine; IL = interleukin; LEF = leflunomide; MTX = methotrexate; NSAID = nonsteroidal anti-inflammatory drug; OSM = oral small molecule; SSZ = sulfasalazine; TNFi = tumor necrosis factor-alpha inhibitor.  
 Singh JA, et al. *Arthritis Rheumatol.* 2019;71:5-32.

14

## Methotrexate Is Not a DMARD in PsA

- 6-month DBRCT of MTX 15 mg/week vs PBO
- Primary outcome: PsARC  
Secondary outcomes: ACR20, DAS28, global and skin scores
- No difference in SJC, TJC, CRP/ESR, PsARC, ACR20, DAS28 at 3 and 6 months
- Patient, MD global, and skin scores significantly improved at 6 months ( $P=0.01, 0.02, 0.02$ )

Measure	Log OR (approx.)	95% CI (approx.)
PsARC	1.5	0.8 - 2.5
ACR20	1.8	1.0 - 3.0
DAS28	1.2	0.7 - 2.0

Despite issues with study design, MTX does not have disease-remitting properties.

CRP = C-reactive protein; DBRCT = double-blind, randomized controlled trial; ESR = erythrocyte sedimentation rate; MTX = methotrexate; PBO = placebo; PsARC = PsA response criteria; DAS = Disease Activity Score; SJC = swollen joint count; TJC = tender joint count.  
 Kingsley GH, et al. *Rheumatology (Oxford).* 2012;51:1368-1377.

15

## CSA in Psoriasis and PsA

- CSA 2.5-5 mg/kg/day yielded PASI75 response in 28% to 97% of patients
- Remission could be maintained at CSA dose of at least 3mg/kg/day
- More than 50% of patients treated with CSA may have an increase in serum creatinine value >30% of baseline if treatment is prolonged for 2 years

### 24 Week Randomized Open NSAID\* Controlled Study of Cyclosporin A in PsA (N=99)

	P-value	Significance CSA vs NSAID*
ACR50	0.02	+
ACR70	0.05	+
Swollen Joint Count	0.05	+
Tender Joint Count	0.01	+
Pain	0.002	+
Patient Global improved ≥1 point	0.04	+
MD Global improved ≥1 point	0.01	+


\*NSAID +/- prednisone 5 mg daily +/- analgesics

Salvanarani C, et al. *J Rheum.* 2001;28:2274-2282. Maza J-H, et al. *JEADV.* 2011;25(2):19-27.

16

## Adalimumab Or Cyclosporine as Monotherapy or Combination For Severe PsA: A Prospective, 12-month, Observational Study


- A 12-month, observational study of 170 TNFi- and cyclosporine-naïve patients
- Patients who received adalimumab (40mg Q2W) (n=57), cyclosporine (2.5-3.75 mg/kg/day) (n=58), or their combination (n=55)
- MTX-IR (25 mg weekly or less, for a minimum of 6 months)
- Assessments: 0.5, 1, 2, 3, 4, 5, 6, 8, 10, and 12 months
- Combination therapy improved PASI50 response rates but NOT beyond the effect of cyclosporine monotherapy (not shown)



MTX-IR = methotrexate inadequate response  
 Karanikolas GN, et al. *J Rheumatol.* 2011;38:2466-2474.

17


# Current and Novel Treatment Options for PsA Treatment



Adapted from Nestle FO, et al. *N Engl J Med.* 2009;361:496-509. Kopf M, et al. *Nat Rev Drug Discov.* 2010;9:703-718. Garber K. *Nat Biotechnol.* 2011;29:563-566. Thanks to Iain McInnes.


18

## TNFi Response in Psoriatic Arthritis in 12 Week Trials (Placebo Corrected)



Adal = adalimumab; Certol = certolizumab; Etan = etanercept; Golim = golimumab; Infix = infliximab.  
Mease PJ, et al. *Rheum Dis N Am.* 2015;4:723-738.

19



20


### Anti-TNFs in PsA: Additional Outcomes

- Enthesitis: ~60–75% improvement
  - Assessment methods evolving: 4-point, MASES, Leeds, SPARCC
- Dactylitis: ~60% improvement
  - Assessment methods evolving: count, score, Leeds dactylometer
- Function
  - Significant improvement achieved as assessed by HAQ
- QoL
  - Significant improvements in SF-36, PsAQoL, DLQI, EQ-5D
- Fatigue
  - Significant improvement observed (eg, FACIT)

MASES = Maastricht Enthesitis Index; SPARCC = Spondyloarthritis Research Consortium of Canada; PsAQoL = PsA quality of life; DLQI = Dermatology Life Quality Index; EQ-5D = EuroQol 5-domain; FACIT = Functional Assessment of Chronic Illness Therapy.  
Mease PJ. *Ann Rheum Dis*. 2011;70(1):i77-i84. Mease PJ. *Arthritis Care Res (Hoboken)*. 2011;63(11):S64-S85.

21

# Current and Novel Treatment Options for PsA Treatment




Adapted from Nestle FO, et al. *N Engl J Med.* 2009;361:496-509. Kopf M, et al. *Nat Rev Drug Discov.* 2010;9:703-718. Garber K. *Nat Biotechnol.* 2011;29:563-566. Thanks to Iain McInnes.

22

## Secukinumab in PsA

### ACR20: Primary Outcome Measure



\*P < 0.0001; †P < 0.001; §P < 0.01; ¶P < 0.05 vs. placebo (P-values at Week 24 adjusted for multiplicity). Missing values imputed as nonresponse (nonresponder imputation). IV = intravenous; SC = subcutaneous.


Mease PJ, et al. *N Engl J Med.* 2015;373:1329-1339. McInnes IB, et al. *Lancet.* 2015;386:1137-1146.

23



## Adalimumab vs Secukinumab in PsA: Indirect Comparison

- N = 302 patients from ADEPT (ADA) and 2 PsA trials of SEC (FUTURE 1 & 2)
- Matching for age, weight, gender, race, MTX use, PASI, dactylitis, enthesitis, HAQ-DI
- Number needed to treat to achieve 1 additional PASI75 responder:
  - ADA 40 mg: 1.7
  - SEC 150 mg: 2.2
  - SEC 300 mg: 1.9




**Conclusion: Secukinumab (anti-IL-17) was shown to be as effective or slightly less effective than adalimumab for PsA (numerical only).**

ADA = adalimumab; SEC = secukinumab.

Betts KA, et al. *Arthritis Rheumatol.* 2015;67(10): Abstract 2868. Strand V. *Rheumatol Ther.* 2017;4:349-362.

24

## CLEAR Study: Secukinumab (aIL-17A) vs Ustekinumab (aIL-12/23) in Psoriasis



Missing data were imputed as nonresponse; only response-evaluable patients were included.

\*P<0.0001; \*\*P=0.0001; †P<0.001; ‡P<0.05

Thaci D, et al. *JAAD.* 2015;73(3):400-409.

25

## Secukinumab: Adverse Events

Common Adverse Events <sup>1</sup>			
	SEC 300 mg	SEC 150 mg	Placebo
URTI	4 (4%)	8 (8%)	7 (7%)
Nasopharyngitis	6 (6%)	4 (4%)	8 (8%)
Diarrhea	2 (2%)	2 (2%)	3 (3%)
Headache	7 (7%)	4 (4%)	4 (4%)
Nausea	3 (3%)	4 (4%)	4 (4%)
Sinusitis	1 (1%)	2 (2%)	1 (1%)
Psoriatic arthropathy	0	3 (3%)	2 (2%)
Urinary tract infection	2 (2%)	4 (4%)	4 (4%)
Hematuria	2 (2%)	3 (3%)	1 (1%)
Vomiting	2 (2%)	2 (2%)	1 (1%)

### Warnings<sup>2</sup>


1. Infection
2. Tuberculosis
3. Hypersensitivity reactions
4. New or worsening inflammatory bowel disease

URTI = upper respiratory tract infection.

1. McInnes IB, et al. *Lancet*. 2015;386:1137-1146. 2. Secukinumab (Cosentyx®) PI 2017 ([www.pharma.us.novartis.com/sites/www.pharma.us.novartis.com/files/cosentyx.pdf](http://www.pharma.us.novartis.com/sites/www.pharma.us.novartis.com/files/cosentyx.pdf)).

26

## SPIRIT-P2: Ixekizumab in Patients with Active PsA and an Inadequate Response to TNFi



Both the 2-week and 4-week ixekizumab dosing regimens improved the signs and symptoms of patients with active PsA who had an inadequate response to TNFi therapy.

\* $P < 0.0001$  vs placebo


IXE = ixekizumab.

Nash P, et al. *Lancet*. 2017;389:2317-2327.

27

## Ixekizumab vs Adalimumab for PsA

- Randomized, double-blind placebo-controlled trial in patients who were biologic DMARD naïve
- More patients achieved an ACR20 response with IXE Q2W (62.1%) or IXE Q4W (57.9%) than placebo (30.2%)
- Disease activity and functional disability were significantly improved with ixekizumab vs placebo ( $P < .01$ ), and there was significantly less progression of structural damage at week 24 with ixekizumab ( $P < .01$ )




**Conclusion: Ixekizumab and adalimumab were both equally better than placebo in PsA. Ixekizumab was better than adalimumab for psoriasis.**

Mease P, et al. *Ann Rheum Dis.* 2017;76:79-87.

28

## SPiRiT H2H: Head-to-Head Comparison of Ixekizumab and Adalimumab



- Ixekizumab was superior to adalimumab in achievement of simultaneous improvement in joint and skin disease (ACR50 and PASI 100) in patients with active PsA and inadequate response to csDMARDs
- Ixekizumab was non-inferior to adalimumab for ACR50 response (IXE: 51%, ADA: 47%) but superior for PASI 100 response (IXE: 60%, ADA: 47%,  $P = .001$ )

Mease PJ, et al. *Ann Rheum Dis.* 2020;79:123-131.

csDMARD = conventional synthetic DMARD

29

## Ixekizumab: Adverse Events

Ixekizumab Adverse Events		
	IXE 80 mg (n=1167)	Placebo (n=791)
Injection site reactions	196 (17%)	26 (3%)
Upper respiratory tract infections	163 (14%)	101 (13%)
Nausea	23 (2%)	5 (1%)
Tinea infections	17 (2%)	1 (<1%)

Adverse events occurring in  $\geq 1\%$  of IXE group, and more frequently than placebo.


### Warnings

1. Infection
2. Tuberculosis
3. Hypersensitivity reactions
4. Inflammatory bowel disease

Ixekizumab (Taltz®) prescribing information (<https://pi.lilly.com/us/taltz-uspi.pdf>)


30

## Current and Novel Treatment Options for PsA Treatment




Adapted from Nestle FO et al. *N Engl J Med.* 2009;361:496-509. Kopf M et al. *Nat Rev Drug Discov.* 2010;9:703-718. Garber K. *Nat Biotechnol.* 2011;29:563-566. Thanks to Iain McInnes.

31



32



33

## Ustekinumab Adverse Events

- Meta-analysis of 9626 patients in 30 RCT of 16 weeks duration
- AEs and SAEs include infections, cough, headache, upper respiratory tract infection, nausea, injection site reactions, CV event, cancer, and death


Adverse events	UST	Placebo	P value
Infections	1210 (19.7%)	588 (17.1%)	<0.01
Nasopharyngitis	318 (5.2%)	162 (4.7%)	0.31
Cough	21 (2.3%)	25 (4.8%)	0.01
Upper respiratory tract infection	150 (3.2%)	201 (7.1%)	<0.001
Nausea	113 (4.8%)	58 (5.0%)	0.80
Headache	302 (6.1%)	141 (5.1%)	0.06
Infusion/Injection site reaction	149 (3.9%)	44 (2.0%)	<0.001
Malignancy	3 (0.1%)	5 (0.2%)	0.16
Death	5 (0.1%)	1 (0.1%)	0.43
CV	7 (0.2%)	4 (0.2%)	1.00

Rolston VS, et al. *Dig Dis Sci.* 2020. doi:10.1007/s10620-020-06344-w.

34

## DISCOVER-1 Trial of Guselkumab

- Phase 3, placebo-controlled trial of guselkumab in 381 patients with active PSA who were biologic-naïve or had previously received a TNFi
- 31% of patients had been previously treated with  $\leq 2$  TNFi agents



	Guselkumab 100 mg		Placebo
	Q4W	Q8W	
Number of patients	128	127	126
ACR20 response at week 24, n/N (%)			
<b>Patients with prior TNFi use</b>	22/38 (57.9%)	23/41 (56.1%)	7/39 (17.9%)
% difference vs. placebo (95% CI)	40.0 (20.8, 59.2)	38.5 (19.3, 57.7)	
Unadjusted p value	<0.001	<0.001	
<b>Patients with inadequate response to prior TNFi</b>	11/17 (64.7%)	9/15 (60.0%)	3/12 (25.0%)
% difference vs. placebo (95% CI)	42.4 (11.0, 73.9)	35.9 (0.8, 71.0)	
<b>Patients without prior TNFi use</b>	54/90 (60.0%)	43/86 (50.0%)	21/87 (24.1%)
% difference vs. placebo (95% CI)	35.9 (22.3, 49.4)	25.9 (12.0, 39.7)	
Unadjusted p value	<0.001	<0.001	


- ACR20 at week 24 was achieved by significantly greater proportions of patients in the guselkumab Q4W (59%) and Q8W (52%) groups than in the placebo group (22%)

Deodhar A, et al. *Lancet.* 2020;395:1115-1125.

35

## DISCOVER-2 Trial of Guselkumab

- Phase 3 trial of 741 biologic-naïve patients with active PsA
- Patients randomized to guselkumab Q4W, Q8W, or placebo



- Significantly greater proportions of patients in the guselkumab Q4W (64%) and Q8W (64%) groups achieved ACR20 at week 24 than placebo (33%)

Mease PJ, et al. *Lancet*. 2020;395:1126-1136.

36


## Guselkumab Adverse Events

	PBO	GUS	
		100 mg Q8W	100 mg Q4W
Patients with $\geq 1$ AE (%)	60%	54%	55%
SAE (%)	4%	3%	0%
Discontinuation due to AE (%)	2%	2%	1%
Infections (%)	25%	26%	24%
Alanine aminotransferase increase	2%	6%	4%
Aspartate aminotransferase increase	2%	7%	2%
Nasopharyngitis	6%	13%	5%
Upper respiratory tract infection	6%	6%	9%

Deodhar A, et al. *Lancet*. 2020;395:1115-1125.

37

# Current and Novel Treatment Options for PsA Treatment




Adapted from Nestle FO et al. *N Engl J Med.* 2009;361:496-509. Kopf M et al. *Nat Rev Drug Discov.* 2010;9:703-718. Garber K. *Nat Biotechnol.* 2011;29:563-566. Thanks to Iain McInnes.

38

## Efficacy of Tofacitinib in PsA

- 395 patients with active PsA and an inadequate response to TNFi were randomized to:
  - Tofacitinib 5 mg BID
  - Tofacitinib 10 mg BID
  - Placebo, with a switch to 5 mg or 10 mg tofacitinib BID at 3 months
- No efficacy noted on Leeds Enthesitis Index, Dactylitis Severity Score, FACIT-F total score, and SF-36 physical functioning

**Conclusion: Tofacitinib has some efficacy in PsA, but no efficacy noted in some symptoms**




Gladman D, et al. *N Engl J Med.* 2017;377:1525-1536.

39



## Adverse Events in 3118 Patient-Years in Tofacitinib Open-Label, Long-Term Extension Study of Therapy for RA\*




\*No dose breakdown; 3227 pts in Treatment Emergent AEs

GI = gastrointestinal disorders; MSK = musculoskeletal and connective tissue disorders; Inf = infections; HGB = decreased hemoglobin; AST/ALT = aspartate/alanine; ANC = absolute neutrophil count.

Wollenhaupt J, et al. ACR 2011. Abstract 407.

40


## Incident Rates of Herpes Zoster in RA Patients



Strangfeld A, et al. EULAR 2020. Abstract OP0238.

41

# Current and Novel Treatment Options for PsA Treatment




Adapted from Nestle FO et al. *N Engl J Med.* 2009;361:496-509. Kopf M et al. *Nat Rev Drug Discov.* 2010;9:703-718. Garber K. *Nat Biotechnol.* 2011;29:563-566. Thanks to Iain McInnes.

42

## Apremilast in PsA: PALACE 1, 2, and 3

Primary endpoint across studies: ACR20 response at week 16




\*P<0.05; §P<0.005; ‡P<0.0001 vs placebo.  
NRI = non-responder imputation

Kavanaugh A, et al. *Ann Rheum Dis.* 2014;73:1020-1026. Cutolo M, et al. *J Rheumatol.* 2016;43:1724-1734. Edwards CJ, et al. *Ann Rheum Dis.* 2016;75:1065-1073.

43


## Apremilast in Moderate-to-Severe Psoriasis

**ESTEEM 1: PASI75 by prior treatment at week 16 (LOCF, full analysis set; N = 844)**



\* $P < 0.0001$ ; <sup>†</sup> $P = 0.0273$  vs PBO; <sup>a</sup>Conventional ± biologics

**ESTEEM 2: PASI75 by prior treatment at Week 16 (LOCF, full analysis set; N = 411)**



\* $P < 0.0001$ ; <sup>†</sup> $P < 0.001$ ; <sup>‡</sup> $P = 0.0069$  vs PBO


LOCF = last observation carried forward.

Paul C et al. *J Am Acad Dermatol.* 2014;70(5):AB164 (abstract P8412). Papp K et al. *J Am Acad Dermatol.* 2015;73:37-49. Reich K et al. AAD 2013, Late breaker. Paul C et al. *Br J Dermatol.* 2015;173:1387-1399.

44

## Apremilast Effects on Enthesitis and Dactylitis

Data pooled from PALACE 1–3, week 24



Gladman DD, et al. *Arthritis Rheum.* 2013;65(10 suppl): S347 (abstract 816).

45

## PALACE 2: 52-Week Safety of Apremilast<sup>1</sup>

Adverse Events	Placebo (N = 159)	APR 30 BID (24 weeks) (N = 162)	APR 30 BID (52 weeks) (N = 234)
Diarrhea	8 (5.0)	24 (14.8)	32 (13.7)
Nausea	3 (1.9)	26 (16.0)	32 (13.7)
Headache	7 (4.4)	19 (11.7)	23 (9.8)
URTI	6 (3.8)	11 (6.8)	22 (9.4)
Nasopharyngitis	6 (3.8)	8 (4.9)	10 (4.3)
Hypertension	7 (4.4)	5 (3.1)	13 (5.6)
<b>Laboratory values</b>			
ALT >150 u/L	1/158 (0.6)	2/160 (1.3)	3/230 (1.3)
Creatinine elevation	0/158 (0.0)	1/160 (0.6)	2/230 (0.9)

### Warnings for<sup>2</sup>:


1. Depression and suicidal behavior
2. Weight loss

APR = apremilast; ALT = alanine aminotransferase.

1. Cutolo M, et al. *J Rheumatol.* 2016;43:1724-1734. 2. Apremilast (Otezla<sup>®</sup>) PI (<http://media.celgene.com/content/uploads/otezla-pi.pdf>).


46

## Current and Novel Treatment Options for PsA Treatment



Adapted from Nestle FO et al. *N Engl J Med.* 2009;361:496-509. Kopf M et al. *Nat Rev Drug Discov.* 2010;9:703-718. Garber K. *Nat Biotechnol.* 2011;29:563-566. Thanks to Iain McInnes.

47



48

## Summary

- Pharmacologic treatment of PsA is only 1 part of the picture. Other factors to consider include:
- Patient goals
  - Improve quality of life, function, and social participation
  - Control symptoms and inflammation (enthesitis, dactylitis, joint pain)
  - Prevent joint damage
- Starting treatment early
- Minimizing associated comorbidities.
- Multidisciplinary care:
  - Physical therapy, occupational therapy, management of comorbidities by dermatologists, endocrinologists, cardiologists, etc.

Perez-Chada LM, et al. *Clin Immunol.* 2020;108397.

49

## Case Study

### Patient with Significant Comorbidities

50

### Case Study: Patient with Significant Comorbidities

- Patient is a 55-year old woman who reports swelling of her left wrist. She complains of lower back pain, bilateral shoulder pain, left wrist and right elbow pain, bilateral 3 PIP and right 3, 4 DIP pain.
  - Patient CDAI: 18 (above TJC and SJC, patient global: 6.0, MD global: 5.0)
  - 2+ edema to mid-calf
- Patient has significant skin involvement (PASI:14)

TJC = Tender Joint Count; SJC = Swollen Joint Count

51

## Lab and Imaging Results

- Lab results:
  - Hemoglobin: **10.0 g/dL** (normal: 12-16)
  - WBC:  $5.2 \times 10^9/L$  (normal: 4.0-11.0)
  - Platelets:  $285 \times 10^9/L$  (normal: 150-400)
  - ESR: **32 mm/hr** (normal: 0-29 mm/hr)
  - Remainder of CBC and CMP are normal.
- Imaging results:
  - Radiographs of the knees shows osteoarthritis on the right.
  - Chest film shows cardiomegaly.

52

## Past Medical History

- Congestive heart failure
- Obesity (BMI: 32)
- Hypertension (160/95 mmHg)
- History of MI three years ago
- Family history positive for MI

*How would you manage this patient?*

53

## Current Visit

- The patient begins taking ixekizumab to control her PsA.
- Two months after starting her therapy, she experiences dyspnea, loss of smell, and a cough for 3 days.
- Her nasal PCR test for COVID-19 is positive.

*How would you manage this patient's PsA given her COVID-19 diagnosis?*

54

## COVID-19 Treatment Modifications

Treatment of Rheumatic Disease During the COVID-19 Pandemic <sup>8</sup>	
Treatment of Rheumatic Disease in the Absence of COVID-19 Infection or Exposure	
HCQ/CQ, SSZ, MTX, LEF, immunosuppressants (tacrolimus, CSA, MMF, AZA), biologics, JAK inhibitors, NSAIDs	Continue therapy
Low-dose corticosteroids	May be started if clinically indicated (<10 mg prednisone equivalent/day)
Following SARS-CoV-2 Exposure	
HCQ/CQ, SSZ, NSAIDs	May be continued
Immunosuppressants (tacrolimus, CSA, MMF, AZA), non-IL-6 biologics, JAK inhibitors	Stop therapy temporarily, pending a negative COVID-19 test or 2 weeks of symptom-free observation
IL-6 inhibitors	May be continued in select circumstances
Documented or presumptive COVID-19	
HCQ/CQ	May be continued
SSZ, MTX, LEF, immunosuppressants, non-IL-6 biologics, and JAK inhibitors	Withhold or stop therapy
NSAIDs	Should be stopped in patients with severe respiratory symptoms

- All recommendations are based on very low quality of evidence and moderate to high consensus.
- The recommendations are for rheumatic disease patients in general and are not subdivided by patient disease. There are no specific recommendations for PsA.
- Mild COVID-19 symptoms: reinitiate therapy in 7-14 days
- Asymptomatic COVID-19: reinitiate therapy in 10-17 days
- Severe COVID-19: reinitiating therapy is dependent on a case-by-case review


Mikuls TR, et al. *Arthritis Rheumatol.* 2020;72:e1-e12.

55



## Conclusions

- A TNFi is recommended as a first-line option in treatment-naïve patients. Contraindications to TNFi therapy include congestive heart failure, previous serious infections, recurrent infections, or demyelinating disease.
- IL-17i or IL-12/23i biologics may be used in patients with severe psoriasis or contraindications to TNFi agents. An IL-17i is recommended over an IL-12/23i, unless the patient has concomitant IBD or prefers less frequent dosing.
- An OSM may be used in patients without severe PsA or severe psoriasis



Singh JA, et al. Arthritis Rheumatol. 2019;71:5-32.

56

## Case Study

### Adverse Events

57

## Initial Presentation

- A 43-year-old overweight woman presents with a 6-year history of PsA principally involving the back. She has difficulty carrying groceries up one flight of stairs due to her back pain. She reports morning stiffness lasting up to 1 hour.
- 
- She also complains of joint pain in her right ankle, right knee, left DIP, and left shoulder.
- Her past medical history is significant for:
  - Type 2 diabetes. HbA1c of 7.6 despite long acting insulin and metformin therapy.
  - Hypertension. Blood pressure of 152/92 mmHg despite lisinopril and furosemide.

58

## Physical Examination

- Patient pain today: 4.5; Patient global today: 5.0; MD global: 5.0
  - Tender bilateral SI joints.
  - **Bilateral Achilles insertional pain.**
  - **Bilateral talocalcaneal pain.**
  - **L23 and 34 pain.**
  - Pain of both shoulders, both wrists, right 3 PIP, bilateral 2 – 4 DIP pain with mild swelling, right knee, right talus pain for a **joint tenderness count of 6/28 and 13/68.**
  - Swelling of right wrist and right knee for a **joint swelling count of 2/28 and 2/66.**  
**CDAI:18.**

59

## Selecting an Initial Treatment Option

- Scaling and mild erythema posterior scalp, thick scaling with mild erythema of both elbows, right intertriginous area and both knees for a **PASI: 8**.

*How would you manage this patient's PsA?*

60

## Initial Treatment

- Teresa is prescribed secukinumab
  - CDAI improved from 18 to 8
  - Mild pain noted in left wrist, DIPs, and right knee only
  - Dactylitis of the right toes
  - PASI improved from 8 to 3
  - Scalp and elbow lesions remain
- Lab results show neutropenia with WBC of 1.4 and PMNs of 0.9

*How would you manage this patient?*

WBC = white blood cell count; PMN = polymorphonuclear cells

61

## Further Management

- Secukinumab therapy is stopped
  - Within 3 weeks, her WBC is 3.2 and PMNs are 2.0
- The patient is started on ustekinumab
  - Patient maintains decrease in CDAI and PASI scores on ustekinumab
  - Lesions on scalp and elbow are improved
  - Patient reports mild tenderness of finger joints

WBC = white blood cell count; PMN = polymorphonuclear cells



62


**Thank You for Your Attention!**



**Med Learning Group - Psoriatic Arthritis**

63


Complimentary poster for the office!  Supplement your Course Learning. It's fast and easy.  We'll ship it to you directly free of charge



## Managing PSORIATIC ARTHRITIS in Specialty Practice:

*New Therapies, Guidelines and Treatment Targets During the COVID-19 Pandemic*

For more information and additional resources please visit [PSORIATICARTHRITIS.POSTERPROGRAM.COM](http://PSORIATICARTHRITIS.POSTERPROGRAM.COM)



64



## Managing PSORIATIC ARTHRITIS in Specialty Practice:

*New Therapies, Guidelines and Treatment Targets During the COVID-19 Pandemic*

[HTTP://WWW.RELIEF-AS.COM](http://WWW.RELIEF-AS.COM)



65