

TeleECHO Series

PRINCIPLES OF TREATMENT:

The Multi-Disciplinary Approach to the Patient
with **Locally Advanced/Metastatic BCC**

Friday, February 12, 2021

FACULTY

Mary L. Stevenson, MD

Assistant Professor

Associate Fellowship Director, Micrographic
Surgery & Dermatologic Oncology


Mohs Micrographic Surgery and Dermatologic Surgery

The Ronald O. Perelman Department of Dermatology

NYU Langone Medical Center

New York, NY

BASAL-CELL CARCINOMA



UMA

COMPLETE
CONFERENCE
MANAGEMENT

UMA



This activity is provided by Med Learning Group.

This activity is co-provided by Ultimate Medical Academy/Complete Conference Management (CCM).

This activity is supported by an independent medical education grant from Regeneron Pharmaceuticals, Inc. and Sanofi Genzyme.



AGENDA

- Basal Cell Carcinoma: Introduction
 - Epidemiology/statistics
 - Prognostic factors
 - Staging
 - NCCN guidelines
 - Considerations for aggressive BCC
- BCC Failing or Not Amenable to Surgery or Radiation
 - Sonic hedgehog pathway
 - Vismodegib and sonidegib
 - Hedgehog side effects
 - Alternate dosing regimens
- When Disease Progresses on Hedgehog Inhibition
 - Immunotherapy and immune checkpoints
 - Cemiplimab, pembrolizumab, etc.
- Case Study Presentation
- Conclusions

***TeleECHO Series: Principles of Treatment:
The Multi-Disciplinary Approach to the Patient With Locally
Advanced/Metastatic BCC***

FACULTY CHAIR

Chrysalyne D. Schmults, MD, MSCE
Associate Professor of Dermatology, Harvard Medical School
Vice-Chair of Surgical Oncology, Brigham and Women's Department of Dermatology
Boston, MA

FACULTY PRESENTERS

Anna Bar, MD

Associate Professor of Dermatology
Co-director of Mohs Micrographic Surgery
Oregon Health & Science University (OHSU)
Portland, OR

Anokhi Jambusaria, MD, MSCE

Assistant Professor, Department of Internal Medicine
Dell Medical School
The University of Texas at Austin
Austin, TX

Justin Leitenberger, MD

Assistant Professor of Dermatology
Oregon Health & Science University (OHSU)
Portland, OR

Emily Ruiz, MD, MPH

Associate Physician, Mohs and
Dermatologic Surgery Center
Dana -Farber/Brigham and Women's Cancer Center
Instructor in Dermatology, Harvard Medical School
Director of the High-Risk Skin Cancer Clinic
Boston, MA

Mary L. Stevenson, MD

Assistant Professor
Associate Fellowship Director, Micrographic
Surgery & Dermatologic Oncology
Mohs Micrographic Surgery and Dermatologic Surgery
The Ronald O. Perelman Department of Dermatology
NYU Langone Medical Center
New York, NY

PROGRAM OVERVIEW

These live virtual TeleECHO® sessions will be a faculty-led didactic and case-based lecture focusing on treatment and management of patients with patients with advanced/metastatic basal cell skin cancer.

TARGET AUDIENCE

This activity is designed to meet the educational needs of medical oncologists, onco-dermatologists, dermatologists, Mohs surgeons, and other healthcare practitioners who care for patients with advanced/metastatic basal cell skin cancer.

LEARNING OBJECTIVES

Upon completion of the program, attendees should be able to:

- Discuss the pathogenesis of metastatic BCC and the attendant involvement of both aberrant pathways and immune dysfunction
- Examine the epidemiology of basal cell carcinoma including prevalence, risk factors, and disease burden
- Apply NCCN practice guidelines for a multi-disciplinary approach to treatment of locally advanced and metastatic BCC
- Explore the clinical trial data for emerging immunotherapy agents in the treatment of advanced or metastatic basal cell skin cancer

ACCREDITATION AND DESIGNATION STATEMENTS

Accreditation Statement

Med Learning Group is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians. This CME activity was planned and produced in accordance with the ACCME Essentials.

Credit Designation Statement

Med Learning Group designates this live virtual activity for a maximum of 1.0 *AMA Category 1 Credit*[™]. Physicians should claim only the credit commensurate with the extent of their participation in the live virtual activity.

Nursing Credit Information

Purpose: This program would be beneficial for nurses involved in the care of patients with advanced/metastatic basal cell skin cancer. **CNE Credits:** 1.0 ANCC Contact Hour.

Accreditation Statement

Ultimate Medical Academy/Complete Conference Management (CCM) is accredited as a provider of continuing nursing education by the American Nurses Credentialing Center's Commission on Accreditation. Awarded 1.0 contact hour of continuing nursing education of RNs and APNs.

ABIM MAINTENANCE OF CERTIFICATION

Successful completion of this CME activity, which includes participation in the evaluation component, enables the participant to earn up to 1.0 Medical Knowledge MOC point in the American Board of Internal Medicine's (ABIM) Maintenance of Certification (MOC) program. It is the CME activity provider's responsibility to submit participant completion information to ACCME for the purpose of granting ABIM MOC credit.

DISCLOSURE POLICY STATEMENT

In accordance with the Accreditation Council for Continuing Medical Education (ACCME) Standards for Commercial Support, educational programs sponsored by Med Learning Group must demonstrate balance, independence, objectivity, and scientific rigor. All faculty, authors, editors, staff, and planning committee members participating in an MLG-sponsored activity are required to disclose any relevant financial interest or other relationship with the manufacturer(s) of any commercial product(s) and/or provider(s) of commercial services that are discussed in an educational activity.

DISCLOSURE OF FINANCIAL RELATIONSHIPS

Chrysalyne D. Schmults, MD, MSCE receives consulting fees from Regeneron Pharmaceuticals and Sanofi. She serves as steering committee member for Castle Biosciences and Regeneron Pharmaceuticals; has received research funding from Castle Biosciences, Regeneron Pharmaceuticals, Novartis, Genentech, and Merck, and is a chair for NCCN.

Anna Bar, MD receives consulting fees from Regeneron Pharmaceuticals. She received research funding to her institution from Castle, Pelle Pharm and Mavis.

Anokhi Jambusaria, MD, MSCE serves on the advisory board for Regeneron Pharmaceuticals.

Justin Leitenberger, MD has nothing to disclose.

Emily Ruiz, MD, MPH receives consulting fees from Sanofi, Pelle Pharm Inc., and Jounce Therapeutic. She discloses other commercial interests with Leo Pharma, Checkpoint Therapeutics, and Jounce Therapeutic.

Mary L. Stevenson, MD states she is a co-investigator for a Regeneron Pharmaceuticals, Inc. clinical trial and receives consulting fees from Two Sigma

CME Content Review

The content of this activity was independently peer reviewed.

The reviewer of this activity has nothing to disclose.

CNE Content Review

The content of this activity was peer reviewed by a nurse reviewer.

The reviewer of this activity has nothing to disclose.

The staff, planners and managers reported the following financial relationships or relationships to products or devices they or their spouse/life partner have with commercial interests related to the content of this CME/CE activity:

Staff, Planners and Managers

Matthew Frese, MBA, General Manager of Med Learning Group, has nothing to disclose.

Christina Gallo, SVP, Educational Development for Med Learning Group, has nothing to disclose.

Chris Drury, Medical Director for Med Learning Group, has nothing to disclose.

Felecia Beachum, Program Manager for Med Learning Group, has nothing to disclose.

Lauren Welch, MA, VP, Outcomes and Accreditation for Med Learning Group, has nothing to disclose.

Brianna Hanson, Outcomes and Accreditation Coordinator for Med Learning Group, has nothing to disclose.

DISCLOSURE OF UNLABELED USE

Med Learning Group requires that faculty participating in any CME activity disclose to the audience when discussing any unlabeled or investigational use of any commercial product or device not yet approved for use in the United States.

During this lecture, the faculty may mention the use of medications for both FDA-approved and nonapproved indications.

METHOD OF PARTICIPATION

There are no fees for participating and receiving CME/CNE credit for this live virtual activity.

To receive CME/CNE credit participants must:

1. Read the CME/CNE information and faculty disclosures.
2. Participate in the live activity.
3. Submit the evaluation form to the Med Learning Group.

You will receive your certificate as a downloadable file.

DISCLAIMER

Med Learning Group makes every effort to develop CME activities that are science-based. This activity is designed for educational purposes. Participants have a responsibility to use this information to enhance their professional development in an effort to improve patient outcomes. Conclusions drawn by the participants should be derived from careful consideration of all available scientific information. The participant should use his/her clinical judgment, knowledge, experience, and diagnostic decision-making before applying any information, whether provided here or by others, for any professional use.

For CME questions, please contact Med Learning Group at info@medlearninggroup.com

Contact this CME provider at Med Learning Group for privacy and confidentiality policy statement information at www.medlearninggroup.com/privacy-policy/

AMERICANS WITH DISABILITIES ACT

Staff will be glad to assist you with any special needs. Please contact Med Learning Group prior to participating at info@medlearninggroup.com



Provided by Med Learning Group



Co-provided by Ultimate Medical Academy/Complete Conference Management (CCM)

This activity is supported by an independent medical education grant from
Regeneron Pharmaceuticals, Inc. and Sanofi Genzyme.

Copyright © 2021 Med Learning Group. All rights reserved. These materials may be used for personal use only. Any rebroadcast, distribution, or reuse of this presentation or any part of it in any form for other than personal use without the express written permission of Med Learning Group is prohibited.

Posting Questions in Zoom Chat

- If you would like to post a question or answer during the presentation, please submit your question or response in the chat feature.
- Remember to direct all questions to the “co-host.” There is a toggle button above the typing space that allows you to specify the location of your message delivery.

TeleECHO Series: Principles of Treatment: The Multidisciplinary Approach to the Patient With Locally Advanced/Metastatic BCC

Mary L. Stevenson, MD

Assistant Fellowship Director, Micrographic Surgery & Dermatologic Oncology
Mohs Micrographic Surgery and Dermatologic Surgery
The Ronald O. Perelman Department of Dermatology
NYU Langone Medical Center
New York, NY

Disclosures

- **Dr. Stevenson** states she is a co-investigator for a Regeneron Pharmaceuticals, Inc. clinical trial and receives consulting fees from Two Sigma.
- During the course of this lecture, Dr. Stevenson may mention the use of medications for both US Food and Drug Administration (FDA)-approved and non-FDA-approved indications
- Case study graphics are not to be copied, reproduced, or distributed in any way

This activity is supported by an independent medical education grant from Regeneron Pharmaceuticals, Inc. and Sanofi Genzyme.

Learning Objectives

- Discuss the pathogenesis of metastatic BCC and the attendant involvement of both aberrant pathways and immune dysfunction
- Examine the epidemiology of BCC including prevalence, risk factors, and disease burden
- Apply practice guidelines for a multidisciplinary approach to the treatment of locally advanced and metastatic BCC
- Explore the clinical trial data for emerging immunotherapy agents in the treatment of advanced or metastatic basal cell skin cancer

Aggressive BCC

Is there such a thing?

Does BCC have a Risk of Metastasis?

- Rates of .0028% to 0.55% reported¹⁻³
- In 1 study of predictive factors: 11,905 BCCs diagnosed between 2000 and 2009 at Brigham and Women's Hospital and Massachusetts General were screened³
- 248 cases $\geq$ 2 cm assessed for risk factors and metastasis/death³
- 248 cases $<$ 2 cm randomly selected for comparison³

1. von Domarus H, Stevens PJ. *J Am Acad Dermatol*. 1984;10(6): 1043-1060. 2. Snow SN, et al. *Cancer*. 1994;73(2):328-335. 3. Morgan F, et al. *J Am Acad Dermatol*. 2020;83:832-838.

Factors predictive of recurrence, metastasis, and death from primary basal cell carcinoma 2 cm or larger in diameter

Frederick C. Morgan, BSPH,^a Emily Stamell Ruiz, MD, MPH,^a Pritesh S. Karia, MPH,^{a,b} Robert J. Besaw, MPH,^a Victor A. Neel, MD, PhD,^c and Chrysalyn D. Schmultz, MD, MSCE
Boston, Massachusetts; Baltimore, Maryland; and Providence, Rhode Island

Multivariable logistic regression of ≥ 2 cm BCC tumors

Variable	Local Recurrence		Metastasis and/or Death	
	OR (95% CI)	P	OR (95% CI)	P
Location Other Head and neck	1 (reference) 9.7 (3.0-31.3)		1 (reference) 5.3 (1.2-23.2)	.026
Diameter Other ≥ 4 cm	— —	—	1 (reference) 11.9 (2.4-59.4)	.003
Tumor depth Other Beyond fat	1 (reference) 3.1 (1.0-9.6)	.049	1 (reference) 28.6 (6.7-121.0)	<.001
Treatment modality Other Mohs micrographic surgery	1 (reference) 0.14 (.04-0.5)	.002	— —	—

BCC > 4 cm on the head and neck, and invading beyond subcutaneous fat, may warrant more aggressive follow-up and radiologic imaging.

Morgan F, et al. *J Am Acad Dermatol.* 2020;83:832-838.

BCC Prognostic Factors

- < 2 cm group had no metastasis/death
- In ≥ 2 cm group, 3 significant predictors of metastasis/death:
 - Diameter ≥ 4 cm or larger
 - Depth beyond subcutaneous fat
 - Head/neck location

Morgan F, et al. *J Am Acad Dermatol.* 2019;83:832-838.

Staging BCC

Tumor Staging System	Definition
AJCC 8th Edition T Staging for Cutaneous Carcinoma of the Head and Neck	
T1	< 2 cm in greatest diameter
T2	≥ 2 cm but < 4 cm in greatest diameter
T3	≥ 4 cm in greatest diameter or minor bone invasion or perineural invasion or deep invasion
T4a	Tumor with gross cortical bone and/or marrow invasion
T4b	Tumor with skull bone invasion and/or skull base foramen involvement
BWH T Staging for BCC	
T1	Tumor diameter < 2 cm or tumor diameter ≥ 2 cm with 0-1 risk factors
T2	Tumor diameter ≥ 2 cm with 2-3 risk factors

Morgan F, Schmultz C, et al. JAAD, in press.

Staging BCC

- BWH T2 BCC:


- Local recurrence: 47% (28%-70%)

- Metastasis/death: 37% (21%-60%)

BWH T2:

≥ 4 cm + depth beyond fat
 ≥ 4 cm + head/neck
 ≥ 2 cm + depth beyond fat + head/neck

Cumulative Incidence of Metastasis or Death by Staging



Morgan F, Schmultz C, et al. JAAD, in press.


CCPDMA Cuts Recurrence Risk in Half for High-Risk BCC

	Recurrence Risk: Recurrent BCC, n (%)	Recurrence Risk: Aggressive Subtype BCC, n (%)
CCPDMA	132/2911 (4.5%)	97/3149 (3%)
SA	32/263 (12%)	17/234 (7%)
p value	p < .001	P < .001
Risk Ratio	.37 [.26-.54]	.42 [.26-.70]

CCPDMA = complete circumferential peripheral and deep margin assessment; SA = Standard Margin Assessment (vertical sections)

Fraga S, Waldman A. Complete marginal assessment in surgically excised keratinocyte carcinoma: A systematic review. Under review.

NCCN: Algorithm for High-Risk BCC



Schmults C, et al. NCCN Guidelines Version 1.2020: Basal Cell Skin Cancer. Available at: https://www.nccn.org/professionals/physician_gls/pdf/nmsc.pdf. Please see NCCN page above for footnotes and additional information/context for these guidelines.

Lack of Adherence to NCCN for High-Risk BCC/SCC

- Survey study of physicians from the 23 NCCN institutions
- ~50% response, N = 57
- Use of CCPDMA in a majority of NCCN high-risk cases
 - 14/15 Mohs surgeons
 - 10/16 other surgeons, dermatologists, radiation oncologists
 - 2/6 pathologists
- Reasons for not using CCPDMA
 - Surgeons defer to pathologists to determine appropriate margin assessment methods
 - Logistical difficulties (time-consuming, overnight stays, open wounds)

NCCN = National Comprehensive Cancer Network; SCC = squamous cell carcinoma.
Danesh M, et al. *Dermatologic Surgery*. 2020;46:1473-1480.

Conclusions: Aggressive BCC

- BWH T2 BCC may have a risk of local recurrence and metastasis in excess of 20%
 - ≥ 4 cm + depth beyond fat
 - ≥ 4 cm + head/neck
 - ≥ 2 cm + depth beyond fat + head/neck
- Do Mohs surgery (or CCPDMA if needed, OR resection)
- Beware of multiply recurrent BCC
- Radiologic surveillance for recurrence
- Consider off-label adjuvant systemic therapy in extreme or multiply recurrent cases

8

Vismodegib

- Indications
 - Metastatic BCC
 - Locally advanced BCC that has recurred following surgery, or in those who are not candidates for surgery/radiation
- Dosage: oral; 150 mg once daily
- Contraindications: none
- Boxed warning: embryo-fetal toxicity


	mBCC (n = 33)	laBCC (n = 63)
ORR	30.3%	42.9%
Complete Response	0%	20.6%
Partial Response	30.3%	22.2%
Median Response Duration	7.6 months	7.6 months

mBCC = metastatic BCC; laBCC = locally advanced BCC; ORR = objective response rate.

Sekulic A, et al. *N Eng J Med*. 2012;366:2171-2179. Vismodegib (Erivedge®) PI 2020 (https://www.gene.com/download/pdf/erivedge_prescribing.pdf). Accessed December 10, 2020.

Vismodegib: ERIVANCE BCC Trial


Progression Free Survival (PFS)



No. at risk

Metastatic	33	31	26	16	12	10	7	6	4	4	3	1	1	0	0	0	0
Locally	63	55	42	38	24	19	17	16	13	10	7	5	3	3	1	1	0
Advanced																	

Overall Survival (OS)



33	33	33	29	25	22	20	18	18	16	15	12	10	7	2	0	0	0
63	61	58	56	52	48	45	43	42	41	40	38	37	25	13	8	2	0

Sekulic A, et al. *BMC Cancer*. 2017;17:332.


Sonidegib

- Indications
 - Locally advanced BCC that has recurred following surgery, or in those who are not candidates for surgery/radiation
- Dosage: oral; 200 mg once daily
- Contraindications: none
- Boxed warning: embryo-fetal toxicity

	laBCC (n = 66)
ORR	56.0%
Complete Response	5%
Partial Response	52%
Median Response Duration	26.1 months

Sonidegib (Odomzo®) PI 2019 (<https://www.odomzo.com/themes/custom/odomzo/global/pdfs/pi.pdf>).

Sonidegib: BOLT Study



BOLT = Basal Cell Carcinoma Outcomes With LDE225 Treatment.
Lear JT, et al. *J Eur Acad Dermatol Venereol*. 2018;32(3):372-381.

Efficacy, safety, and comparison of sonic hedgehog inhibitors in basal cell carcinomas: A systematic review and meta-analysis

Pingxing Xie, MD, PhD, and Philippe Lefrançois, MD, PhD
Montreal, Canada

- 16 studies: quantitative meta-analysis of safety and efficacy
- Locally advanced BCC
 - Overall Response Rate: comparable for vismodegib and sonidegib (69% vs 57%)
 - Complete Response Rate: superior for vismodegib (31% vs 3%)
- Metastatic BCC
 - Overall Response Rate: superior for vismodegib (39% vs 15%)
- Side effects (combined prevalence)
 - 67%, 54%, and 58% for muscle spasms, dysgeusia, and alopecia, respectively; comparable for sonidegib and vismodegib
 - Upper GI distress more common in sonidegib than in vismodegib use
- Vismodegib favored over sonidegib in clinical practice

Xie P, Lefrançois P. *J Am Acad Dermatol* 2018;79:1089-1100.

Hedgehog Side Effects

- Hedgehog inhibitor side effects are not life-threatening but are nearly universal and often severe
- High rate of discontinuation in clinical trials (20%-50%)
 - Debilitating muscle pain/cramping
 - Loss of taste leading to major weight loss
 - Hair loss is often severe
- Alternatives to daily dosing are often used (though minimally studied)
 - A dosing regimen of 1 to 2 weeks on, 1 week off, is what we employ
 - Impact on efficacy unknown but such modifications are often needed to remain on treatment

Migden M, et al. *Lancet Oncology*. 2015;16:716-728. Dummer R, et al. *Br J Dermatol*. 2020;182(6):1369-1378. Sekulic A, et al. *BMC Cancer*. 2017;17:332. Sekulic A, et al. *N Eng J Med*. 2012;366:2171-2179. Villani A, et al. *Dermatol Ther*. 2020;10:401-412.


Alternate Dosing Regimens

- More tolerable than continuous therapy, with comparable effectiveness
- Impact on secondary resistance undefined
- BCC managed using hedgehog inhibitor (HHI) is comparable to chronic disease management or palliation
- May take months, weeks, or days off without loss of effectiveness



Two intermittent vismodegib dosing regimens in patients with multiple basal-cell carcinomas (MIKIE): a randomised, regimen-controlled, double-blind, phase 2 trial

Brigitte Dréno, Rainer Kunstfeld, Axel Hauschild, Scott Fosko, David Zloty, Bruno Labeille, Jean-Jacques Grob, Susana Puig, Frank Gilberg, Daniel Bergström, Damian R Page, Gary Rogers, Dirk Schadendorf



Dréno B, et al. *Lancet Oncol.* 2017; 18:404-412.

RESEARCH LETTER

A Novel Alternate Dosing of Vismodegib for Treatment of Patients With Advanced Basal Cell Carcinomas

Vismodegib Dosing Regimen, Response, and Adverse Effects

Patient No./Sex	BCC History	IVT Regimen (with/without medication = no. of cycles)	Length of Follow-Up, mo	BCC Response	Adverse Effects
1/M	>50 BCCs	1 wk/1 wk x 8 cycles; 1 wk/3 wks x 12 cycles	20	Decrease in number and size of lesions	Fatigue, arthralgia, arthritis, and alopecia; resolved after decreasing dose
2/M	Basal cell nevus syndrome	1 wk/2 wks x 10 cycles	12	Decrease in number and size of lesions	Fatigue, dysgeusia, and muscle aches
3/F	Multiple BCCs on the face	1 wk/1 wk x 6 cycles; 1 wk/2 wks x 7 cycles; 1 wk/3 wks x 14 cycles	24	Resolution	Alopecia; resolved with decreased dosage
4/M	Multiple recurrent and aggressive BCCs on face	1 wk/1 wk x 8 cycles; 1 wk/2 wks x 1 cycle; 1 wk/3 wks x 9 cycles	19	Resolution	Fatigue, dysgeusia, muscle spasms, and hand arthritis; improved with lowered dosage
5/F	Large periorbital BCC	1 wk/1 wk x 6 cycles; 1 wk/2 wks x 5 cycles	12	Decrease in BCC size	Fatigue, dysgeusia, muscle aches, and alopecia
6/M	Large BCC on nose	1 wk/1 wk x 6 cycles	7	Decrease in BCC size, followed by MMS	Fatigue, dysgeusia, diarrhea, and constipation
7/M	Incompletely excised periorbital BCC	1 wk/3 wks x 5 cycles	9	Resolution	Fatigue

7 patients: 1 week on/1 to 3 weeks off based on tolerability; 3 BCC clinically resolved, 4 decreased in size
No patients discontinued therapy based on adverse effects.

IVT = intermittent vismodegib therapy; MMS = Mohs micrographic surgery.
Becker LR, et al. *JAMA Dermatol.* 2017; 153:321-322.

Outcomes for Basal Cell Carcinoma Treated With Vismodegib Extended Alternate Day Dosing

- 8 patients
- Vismodegib 150 daily x 3 months, then extended alternate day dosing as tolerated
 - Every other day to once weekly dosing
 - Mean reduction in tumor size: ~50%
 - 4 patients experienced no side effects, 4 experienced mild side effects that
 - improved/resolved with further extended alternate day dosing




Before

Every 4 days x 10 months

Routt E, Ratner D. *Dermatol Surg.* 2020;46:1109-1112.

When Disease Progresses on Hedgehog Inhibition?


Immunotherapy: The Cancer-Immunity Cycle



APCs = antigen-presenting cells; CTLs = cytotoxic T lymphocytes.
 Chen DS, Mellman I. *Immunity*. 2013;39:1-10.

Immune Checkpoints


Activation of T lymphocytes




CTLA4 = cytotoxic T lymphocyte-associated protein 4; PD1 = programmed cell death 1; PDL1 = PD1 ligand 1; TIM3 = T cell immunoglobulin and mucin domain-containing protein 3; VISTA = V-domain immunoglobulin suppressor of T cell activation.
Sharma P et al. *Nat Rev Cancer*. 2011;11:805-812.

Immune Checkpoints

Activation of T lymphocytes




Inhibition of T lymphocytes



CTLA4 = cytotoxic T lymphocyte-associated protein 4; PD1 = programmed cell death 1; PDL1 = PD1 ligand 1; TIM3 = T cell immunoglobulin and mucin domain-containing protein 3; VISTA = V-domain immunoglobulin suppressor of T cell activation.
Sharma P et al. *Nat Rev Cancer*. 2011;11:805-812.

Immune Checkpoints



Cemiplimab

- Feb 9, 2021: First FDA-approved immunotherapy for BCC (advanced cases previously treated with HHI)
 - Also indicated for metastatic cutaneous SCC and laSCC

Trial of cemiplimab for those not tolerating or progressing on HHI

	laBCC (n = 84)
ORR	31.0%
Complete Response	6.0%
Partial Response	25.0%
Response	85% responses ongoing at 12 months
Estimated PFS (all patients)	19 months

Stratigos AJ, et al. *Ann Oncology*. 2020;31:S1175-S1176.

Cemiplimab

- Median baseline TMB: 58.2 (responding; n = 18) and 23.5 (non-responding; n = 38) mutations/Mb
- Responses occurred at all TMB levels
- Downregulation of MHC-I expression identified as an immune evasion mechanism in non-responding BCCs with high TMB
- Adverse events: fatigue (30%), diarrhea (24%) and pruritus (21%)
- 17% of patients discontinued due to AEs

TMB = tumor mutational burden; MHC = major histocompatibility complex.
Stratigos AJ, et al. *Ann Oncology*. 2020;31:S1175-S1176.

Pembrolizumab With/Without Vismodegib for Advanced BCC


	All Participants (N = 16)	Pembrolizumab (n = 9)	Pembrolizumab + Vismodegib (n = 7)
ORR	38%	44%	29%
1-yr PFS	70%	62%	83%
1-yr OS	94%	89%	100%

- No life-threatening AEs or deaths during the study
- 3 grade 3 AEs occurred out of 98 AEs from 16 participants (1 case of hyponatremia attributed to pembrolizumab)
- 23 immune-related AEs (grade 1/2 dermatitis and fatigue most common)

Chang A, et al. *J Am Acad Dermatol*. 2019;564-566.

Not currently FDA approved for BCC

Pembrolizumab: Best % Change in BCC Diameter From Baseline




Chang A, et al. *J Am Acad Dermatol.* 2019;564-566.

Not currently FDA approved for BCC

Summary

- Advanced BCC is best managed using multidisciplinary consultation
- Diameter > 4 cm, head/neck location, and/or invasion beyond subcutaneous fat significantly increase risk for local recurrence, metastasis, and death
- Vismodegib outperforms sonidegib in a recent meta-analysis on safety and efficacy
- Alternate dosing regimens increase tolerability without compromising efficacy
- Immune checkpoint therapy is now approved (cemiplimab) and is useful in cases of hedgehog inhibitor progression or intolerance



Cases

Case 1

Locally Aggressive BCC: Case 1

Courtesy of Dr. Chrysalyne Schmults

- A 58-year-old healthy man with BCC on right nasal sidewall initially presented in 2012
- Status post (S/P) radiation as primary therapy “to avoid bone resection near eye”

Aggressive BCC

- 2015 to 2016: 5 subsequent recurrences excised by plastic surgeon with in-office frozen sections read by pathologist as clear
- At sixth recurrence the patient is referred for salvage radiation
- Radiation oncologist, in turn, referred the patient to the cancer center


What would be your next step in assessing this patient?

Aggressive BCC

- Magnetic resonance imaging (MRI): tumor abuts bone, no sinus/orbit/nerve invasion
- No metastasis on neck/chest computed tomography (CT)



Mohs Pathology



The tumor abuts perichondrium/cartilage, infiltrating between septal cartilages.

- Mohs cleared all but nasal bone and turbinate mucosa (purple outline)
- Head/neck surgeon Dr. Jason Kass requested en face assessment of posterior resection



Locally Aggressive BCC: Polling Question

The NCCN recommends which of the following at this point?

1. CCPDMA
2. Radiation
3. Systemic agent
4. Something else


- Mohs cleared all but nasal bone and turbinate mucosa (purple outline)
- Head/neck surgeon Dr Jason Kass requested en face assessment of posterior resection
- **NCCN: Recommends CCPDMA for high-risk BCC that can't be closed primarily**



Margin clear but
bony margin not
processed en face



Septal bone graft
(from radius) and
free flap via
Dr. Jason Kass



- Local recurrence: 47% (28%-70%)
- Metastasis: 37% (21%-60%)
- Adjuvant radiation?
 - No data on risk reduction
 - May compromise current reconstruction and make future surgery difficult
- Adjuvant vismodegib?
 - No data on risk reduction in adjuvant setting
 - Difficult side effect profile
 - Offered but declined
- Follow-up every 6 months with face MRI and neck CT every 2 to 3 years

At 1 year, the patient is doing well.





- At 1.5 years, notes nasal congestion, clotted blood
- Recurrent tumor focus, right ethmoid sinus, superior turbinate
- Recurred where couldn't complete Mohs resection
- Another OR resection planned

Despite Pre-Op Imaging and Multiple Scouting Biopsies to Aid Resection Planning, Margins Were Diffusely Positive

- In bone, underside of prior flap, lacrimal duct
- Patient declined rhinectomy and further maxilla resection

Next Steps for This Patient

- Vismodegib daily with complete response radiologically at 8 months but with extreme cachexia
- Tumor recurred during drug holiday
 - Again declined surgery
 - Currently weighing radiation vs vismodegib retreatment vs cemiplimab

Case 2

BCC With Distant Metastasis

Case courtesy of Dr Emily Ruiz

- 60-year-old male
- 2000: initial treatment of BCC on left upper back with ED&C
- 2011: retreated with ED&C but lesion never healed
- 2013: second recurrence, biopsy-proven infiltrative BCC, treated with excision with frozen sections with negative margins on final sections
- 2017: developed axillary pain that was not worked up
- 2018: incidental lung tumors noted on CT scan performed for another chief complaint
- 2018: biopsy of axillary node confirmed diagnosis of metastatic BCC

ED&C: electrodesiccation and curettage

BCC With Distant Metastasis: Failure

Case courtesy of Dr Emily Ruiz




Locally Aggressive BCC: Polling Question

Which of the following options would you try at this point?

1. Radiation
2. Surgery
3. Hedgehog inhibitor
4. Anti-PD1 inhibitor
5. Conventional chemotherapy
6. Observation




BCC With Distant Metastasis: Failure

Case courtesy of Dr Emily Ruiz







BCC With Distant Metastasis: Failure

Case courtesy of Dr Emily Ruiz



BCC With Distant Metastasis: Failure


Case courtesy of Dr Emily Ruiz



What would you try next for this patient?

BCC With Distant Metastasis: Success



Case courtesy of Dr Emily Ruiz



Before Cemiplimab (vismodegib failure)

BCC With Distant Metastasis: Success


Case courtesy of Dr Emily Ruiz



Conclusions

- BCC is the most common form of skin cancer worldwide
- BCCs are slow-growing and rarely metastasize; however, they are locally invasive and can be destructive
- BCC pathogenesis and the implications of aberrant pathways and immune dysfunction continue to be clarified
- Therapy selection depends on the patient's age and gender, along with site, size, and type of lesion
- Current mainstay of BCC treatment involves surgical modalities (excision, electrodesiccation and curettage [EDC], cryosurgery, Mohs surgery)
- Systemics include the HHIs vismodegib and sonidegib; there is now FDA-approved therapy for laBCC post-HHI (cemiplimab – Feb/2021); other PD1/PDL1 inhibitors are under investigation

Thank You!



Complimentary
poster for the
office!

Supplement your
Course Learning.
It's fast and easy.



We'll ship it
to you directly
free of charge



BASAL-CELL CARCINOMA

TeleECHO Series

PRINCIPLES OF TREATMENT:
The Multi-Disciplinary Approach to the Patient with Locally Advanced/ Metastatic BCC

For more information and additional resources please visit
BCC.POSTERPROGRAM.COM

Basal Cell Carcinoma: Identification and Management

Resource	Address
Ascierto PA, et al. Immunotherapy in non-melanoma skin cancer: Updates and new perspectives. <i>Drugs Context</i> . 2019;8:212583.	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6434981/
Carballo GB, et al. A highlight on sonic hedgehog pathway. <i>Cell Commun Signal</i> . 2018;16:11.	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5861627/
Cameron MC, et al. Basal cell carcinoma: Epidemiology; pathophysiology; clinical and histological subtypes; and disease associations. <i>J Am Acad Dermatol</i> . 2019;80:303-317.	https://pubmed.ncbi.nlm.nih.gov/29782900/
Cameron MC, et al. Basal cell carcinoma: Contemporary approaches to diagnosis, treatment, and prevention. <i>J Am Acad Dermatol</i> . 2019;80:321-339.	https://pubmed.ncbi.nlm.nih.gov/29782901/
Chang ALS, et al. Pembrolizumab for advanced basal cell carcinoma: An investigator-initiated, proof-of-concept study. <i>J Am Acad Dermatol</i> . 2019;80:564-566.	https://www.jaad.org/article/S0190-9622(18)32471-X/pdf
Chen DS, et al. Oncology meets immunology: The cancer-immunity cycle. <i>Immunity</i> . 2013;39:1-10.	https://pubmed.ncbi.nlm.nih.gov/23890059/
Danesh MJ, et al. Adherence to the National Comprehensive Cancer Network Criteria of Complete Circumferential Peripheral and Deep Margin Assessment in Treatment of High-Risk Basal and Squamous Cell Carcinoma. <i>Dermatologic Surg</i> . 2020;46:1473-1480.	https://pubmed.ncbi.nlm.nih.gov/32149872/
Dréno B, et al. Two intermittent vismodegib dosing regimens in patients with multiple basal-cell carcinomas (MIKIE): A randomised, regimen-controlled, double-blind, phase 2 trial. <i>Lancet Oncol</i> . 2017;18:404-412.	https://www.thelancet.com/journals/lanonc/article/PIIS1470-2045(17)30072-4/fulltext

Dummer R, et al. Long-term efficacy and safety of sonidegib in patients with advanced basal cell carcinoma: 42-month analysis of the phase II randomized, double-blind BOLT study. <i>Br J Dermatol</i> . 2020;182:1369-1378.	https://pubmed.ncbi.nlm.nih.gov/31545507/
McDaniel B, et al. Basal Cell Carcinoma. In: StatPearls [Internet]. Treasure Island, FL: StatPearls Publishing. Updated November 20, 2020.	https://www.ncbi.nlm.nih.gov/books/NBK482439/
Migden MR, et al. Emerging trends in the treatment of advanced basal cell carcinoma. <i>Cancer Treat Rev</i> . 2018;64:1-10.	https://pubmed.ncbi.nlm.nih.gov/29407368/
Morgan FC, et al. Factors predictive of recurrence, metastasis, and death from primary basal cell carcinoma 2 cm or larger in diameter. <i>J Am Acad Dermatol</i> . 2020;83:832-838.	https://pubmed.ncbi.nlm.nih.gov/31600531/
Paulson KG, et al. Immunotherapy for skin cancer. <i>Int Immunol</i> . 2019;31:465-475.	https://pubmed.ncbi.nlm.nih.gov/30753483/
Sekulic A, et al. Long-term safety and efficacy of vismodegib in patients with advanced basal cell carcinoma: final update of the pivotal ERIVANCE BCC study. <i>BMC Cancer</i> . 2017;17:332.	https://bmccancer.biomedcentral.com/articles/10.1186/s12885-017-3286-5
Sekulic A, et al. Efficacy and safety of vismodegib in advanced basal-cell carcinoma. <i>N Eng J Med</i> . 2012;366:2171-2179.	https://www.nejm.org/doi/full/10.1056/nejmoa1113713
Stratigos AJ, et al. LBA47 Primary analysis of phase II results for cemiplimab in patients (pts) with locally advanced basal cell carcinoma (laBCC) who progress on or are intolerant to hedgehog inhibitors (HHIs). <i>Ann Oncology</i> . 2020;31(suppl 4):S1175-S1176.	https://www.annalsofoncology.org/article/S0923-7534(20)42359-2/abstract
Stratigos AJ, et al. European interdisciplinary guideline on invasive squamous cell carcinoma of the skin: Part 2. Treatment. <i>Eur J Cancer</i> . 2020;128:83-102.	https://pubmed.ncbi.nlm.nih.gov/32113942/

Tanese K. Diagnosis and management of basal cell carcinoma. <i>Curr Treat Options Oncol</i> 2019;20:13.	https://pubmed.ncbi.nlm.nih.gov/30741348/
Totonchy M, et al. Emerging concepts and recent advances in basal cell carcinoma. <i>F1000Res</i> . 2017;6:2085.	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5717469/
Xie P, et al. Efficacy, safety, and comparison of sonic hedgehog inhibitors in basal cell carcinomas: A systematic review and meta-analysis. <i>J Am Acad Dermatol</i> . 2018;79:1089-1100.e17.	https://pubmed.ncbi.nlm.nih.gov/30003981/

Resources and Societies

Resource	Address
American Academy of Dermatology (AAD). <i>Skin Cancer</i> .	https://www.aad.org/media/stats-skin-cancer
American Cancer Society (ACS). <i>Basal and Squamous Cell Skin Cancer</i> .	https://www.cancer.org/cancer/basal-and-squamous-cell-skin-cancer.html
American Society of Clinical Oncology (ASCO).	https://www.asco.org/
Centers for Disease Control and Prevention (CDC). <i>What Is Skin Cancer?</i>	https://www.cdc.gov/cancer/skin/basic_info/what-is-skin-cancer.htm
National Comprehensive Cancer Network. <i>NCCN Clinical Practice Guidelines in Oncology. Basal Cell Skin Cancer. Version 1.2020</i> .	https://www.nccn.org/professionals/physician_gls/pdf/nmsc.pdf
Skin Cancer Foundation. <i>Basal Cell Carcinoma Overview</i> .	https://www.skincancer.org/skin-cancer-information/basal-cell-carcinoma/