

A NEW LIGHT IN THE DARKNESS:


*New Virus-neutralizing Monoclonal Antibodies and
Other Point-of-Care Therapies Recently Granted Emergency
Use Authorizations for Patients with COVID-19*

MEETING INFO

Saturday, March 27,
2021

FACULTY

Christopher Palma, MD, ScM
Assistant Professor of Medicine
University of Rochester
Rochester, NY



A LIGHT IN THE DARKNESS:

New Virus-neutralizing Monoclonal Antibodies and Other Point-of-Care Therapies Recently Granted Emergency Use Authorizations for Patients with COVID-19

AGENDA

1. **Rationale for the Use of New Virus-neutralizing Monoclonal Antibodies**
 - a. High mutation rate of RNA viruses
 - b. The risk of viral mutations leading to therapy resistance (***Whiteboard theme: depiction of viral mutations leading to therapy resistance***)
 - c. Mechanism of action of new virus-neutralizing monoclonal antibodies in mitigating the risk of viral resistance to therapy

2. **Therapies Granted Emergency Use Authorization for Patients with COVID-19**
 - a. What is emergency use authorization?
 - b. Clinical trial data on the efficacy and safety of new virus-neutralizing monoclonal antibodies and other therapies approved for emergency use in all patients who test positive for COVID-19 (***Whiteboard theme: MOA of new virus-neutralizing monoclonal antibodies approved for emergency use in all patients who test positive for COVID-19***)
 - c. Guidance of the development of in-clinic infusion capability to deliver new virus-neutralizing monoclonal antibodies at the point-of-care

3. **COVID-19 Vaccine Development**
 - a. Efficacy and safety of the first FDA-granted emergency use authorization vaccine for the prevention of COVID-19
 - b. Updates on vaccines in development

4. **Case studies**

5. **Conclusions**

A Light in the Darkness: New Virus-neutralizing Monoclonal Antibodies and Other Point-of-Care Therapies Recently Granted Emergency Use Authorization for Patients with COVID-19

FACULTY

PROGRAM CHAIR

Shyam Kottilil, MD, PhD (PROGRAM CHAIR)

Professor of Medicine
Chief, Division of Infectious Diseases
Institute of Human Virology
University of Maryland
Baltimore, MD

FACULTY PRESENTERS

Roger Bedimo, MD, MS

Professor of Medicine
University of Texas Southwestern Medical Center
Dallas, TX

William A. Fischer II, MD

Associate Professor of Medicine, Pulmonary and Critical Care Medicine
Director of Emerging Pathogens
Institute for Global Health and Infectious Diseases
The University of North Carolina
Chapel Hill, NC

Shivakumar Narayanan, MBBS, MD

Assistant Professor
Division of Clinical Care and Research
Director, Institute of Human Virology
University of Maryland, School of Medicine
Baltimore, MD

Christopher Palma, MD, ScM

Assistant Professor of Medicine
University of Rochester
Rochester, NY

PROGRAM OVERVIEW

The COVID-19 FRONTLINE TeleECHO series provides a comprehensive and up-to-date perspective on the ever-changing management of patients with COVID-19. Each TeleECHO session features in-depth case studies to encourage retention of the lessons and provide new perspectives on the management of patients during the COVID-19 pandemic. The case studies will focus on different issues facing clinicians, such as identifying patients who would benefit from monoclonal antibody therapy and best practices for incorporating agents authorized for emergency use into the care of hospitalized and non-hospitalized patients with COVID-19. Strategies for administering neutralizing monoclonal antibodies, such as referral to local infusion centers or developing in-clinic infusion capabilities, will also be discussed.

TARGET AUDIENCE

This CME initiative is designed for HCPs who are involved in the care and treatment of patients with COVID-19 in an outpatient setting, including physicians, NPs, PAs, nurses, pharmacists and paramedics.

LEARNING OBJECTIVES

Upon the completion of this program, attendees should be able to:

- Assess the rationale for the use of neutralizing monoclonal antibody therapies in recently diagnosed COVID-19 patients to prevent the development of severe disease
- Critique the efficacy and safety of new virus-neutralizing monoclonal antibody therapies and other therapies approved for emergency use in all patients who test positive for COVID-19
- Develop in-clinic infusion capability in order to administer new virus-neutralizing monoclonal antibodies to patients with COVID-19 at the point-of-care

ACCREDITATION STATEMENT

Med Learning Group is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

This CME activity was planned and produced in accordance with the ACCME Essentials.

CREDIT DESIGNATION STATEMENT

Med Learning Group designates this live virtual activity for a maximum of 1.0 *AMA Category 1 Credit*[™].

Physicians should claim only the credit commensurate with the extent of their participation in the live virtual activity.

NURSING CREDIT INFORMATION

Purpose: This program would be beneficial for nurses involved in the treatment of patients with COVID-19.

Credits: 1.0 ANCC Contact Hour.

CNE Accreditation Statement: Ultimate Medical Academy/CCM is accredited as a provider of continuing nursing education by the American Nurses Credentialing Center's Commission on Accreditation. Awarded 1.0 contact hour of continuing nursing education of RNs and APNs.

ABIM Maintenance of Certification

Successful completion of this CME activity, which includes participation in the evaluation component, enables the participant to earn up to 1.0 Medical Knowledge MOC point in the American Board of Internal Medicine's (ABIM) Maintenance of Certification (MOC) program. It is the CME activity provider's responsibility to submit participant completion information to ACCME for the purpose of granting ABIM MOC credit.

DISCLOSURE POLICY STATEMENT

In accordance with the Accreditation Council for Continuing Medical Education (ACCME) Standards for Commercial Support, educational programs sponsored by Med Learning Group must demonstrate balance, independence, objectivity, and scientific rigor. All faculty, authors, editors, staff, and planning committee

members participating in an MLG-sponsored activity are required to disclose any relevant financial interest or other relationship with the manufacturer(s) of any commercial product(s) and/or provider(s) of commercial services that are discussed in an educational activity.

DISCLOSURE OF CONFLICTS OF INTEREST

Faculty Member	Disclosures
Shyama Kottlilil, MD, PhD	Discloses that the University of Maryland has received funds to participate in trials, as well he has received research funds paid to the university from Merck Inc, Gilead Sciences and Arbutus Pharmaceuticals. He has also provided contracted research for Regeneron, Eli Lilly, and air Pharmaceuticals, as well as serving on the advisory board for hepatitis b functional cure program at Merck Inc.
Roger Bedimo, MD, MS	Discloses that he has worked as a Consultant for Merck & Co, Viiv Healthcare and Theratechnologies.
William A. Fischer II, MD	Discloses that he has been contracted for research for Ridgeback Biopharmaceuticals for COVID-19 research, as well as worked as Consulted for Merck and Roche. He also worked for Syneos and Janssen for adjudication of AE in RSV and Influenza studies respectively, and served as the site PI for the Phase I Lilly study of - Bamlanivimab and for the Phase II study of Casirivimab/Imdevimab at University of North Carolina.
Shivakumar Narayanan, MBBS, MD	Discloses that has received contracted research funding from Regeneron Pharmaceuticals. Dr. Narayanan has been an investigator in clinical studies sponsored by Regeneron Pharmaceuticals. He is a principal investigator and a recipient of contracted research funding from Gilead Sciences.
Christopher Palma, MD, ScM	Discloses that he has been contracted for research for Regeneron.

CME Content Review

The content of this activity was independently peer reviewed.

The reviewer of this activity has nothing to disclose.

CNE Content Review

The content of this activity was peer reviewed by a nurse reviewer.

The reviewer of this activity has nothing to disclose.

Staff Planners and Managers

The staff, planners, and managers reported the following financial relationships or relationships to products or devices they or their spouse/life partner have with commercial interests related to the content of this CME/CE activity:

Matthew Frese, MBA, General Manager of Med Learning Group, has nothing to disclose.

Christina Gallo, SVP, Educational Development for Med Learning Group, has nothing to disclose.

Diana Tommasi, PharmD, Medical Director for Med Learning Group, has nothing to disclose.

Lauren Welch, MA, VP, Accreditation and Outcomes for Med Learning Group, has nothing to disclose.

Lisa Crenshaw, Senior Program Manager for Med Learning Group, has nothing to disclose.

Russie Allen, Accreditation and Outcomes Coordinator for Med Learning Group, has nothing to disclose.

Morgan Kravarik, Program Coordinator for Med Learning Group, has nothing to disclose.

Daniel Dasilva, Accreditation and Outcomes Coordinator for Med Learning Group, has nothing to disclose.

DISCLOSURE OF UNLABELED USE

Med Learning Group requires that faculty participating in any CME activity disclose to the audience when discussing any unlabeled or investigational use of any commercial product or device not yet approved for use in the United States.

During this lecture, the faculty may mention the use of medications for both FDA-approved and non-approved indications.

METHOD OF PARTICIPATION

There are no fees for participating and receiving CME credit for this live virtual activity. To receive CME/CNE credit participants must:

1. Read the CME/CNE information and faculty disclosures.
2. Participate in the web-based live activity.
3. Submit the evaluation form to Med Learning Group.

You will receive your certificate upon completion.

DISCLAIMER

Med Learning Group makes every effort to develop CME activities that are science based.

This activity is designed for educational purposes. Participants have a responsibility to use this information to enhance their professional development in an effort to improve patient outcomes. Conclusions drawn by the participants should be derived from careful consideration of all available scientific information. The participant should use his/her clinical judgment, knowledge, experience, and diagnostic decision making before applying any information, whether provided here or by others, for any professional use.

For CME questions, please contact Med Learning Group at info@medlearninggroup.com

Contact this CME provider at Med Learning Group for privacy and confidentiality policy statement information at <http://medlearninggroup.com/privacy-policy/>



Provided by Med Learning Group



Co-provided by Ultimate Medical Academy/Complete Conference Management (CCM).

This activity is supported by an educational grant from Lilly.

Copyright © 2021 Med Learning Group. All rights reserved. These materials may be used for personal use only. Any rebroadcast, distribution, or reuse of this presentation or any part of it in any form for other than personal use without the express written permission of Med Learning Group is prohibited.

COVID-19 FRONTLINE

A Light in the Darkness: New Virus-neutralizing Monoclonal Antibodies and Other Point-of-Care Therapies Recently Granted Emergency Use Authorizations for Patients with COVID-19

Christopher Palma, MD, ScM
Assistant Professor of Medicine
University of Rochester
Rochester, NY

Disclosures

- Dr. Palma discloses that he has been contracted for research for Regeneron.
- During this lecture, Dr. Palma may mention the use of medications for both FDA-approved and nonapproved indications.


This activity is supported by an independent medical education grant from Lilly.

Learning Objectives

- Assess the rationale for the use of neutralizing monoclonal antibody therapies in recently diagnosed COVID-19 patients to prevent the development of severe disease
- Critique the efficacy and safety of new virus-neutralizing monoclonal antibody therapies and other therapies approved for emergency use in all patients who test positive for COVID-19
- Develop in-clinic infusion capability in order to administer new virus-neutralizing monoclonal antibodies to patients with COVID-19 at the point-of-care

SARS-CoV-2

- COVID-19 is caused by the SARS-CoV-2 virus¹⁻³
- The virus is spread primarily via respiratory droplets during face-to-face contact²
- Spike protein on viral surface binds to ACE2 receptor on target cells, facilitating viral entry into host cells^{2,3}




SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; COVID-19 = coronavirus disease 2019; ACE = angiotensin-converting enzyme.

1. Adapted from Vaduganathan M, et al. *N Engl J Med.* 2020;382:1653-1659. 2. Wiersinga WJ, et al. *JAMA.* 324:782-793. 3. Baum A, et al. *Science.* 2020;369:1014-1018.

Clinical Presentation of COVID-19

Systemic and respiratory disorders caused by COVID-19

Systemic Disorders		Respiratory Disorders
Fever, cough, fatigue, sputum production, headache		Rhinorrhea, sneezing, sore throat
Hemoptysis, acute cardiac injury		Pneumonia
Hypoxemia		Ground-glass opacities
Dyspnea, Lymphopenia		RNAemia, acute respiratory distress syndrome
Diarrhea		

Most common symptoms of COVID-19 at presentation

Symptom	Patients Presenting with Symptom (N = 1420)
Headache	70.3%
Loss of smell	70.2%
Nasal obstruction	67.8%
Asthenia	63.3%
Cough	63.2%
Myalgia	62.5%
Rhinorrhea	60.1%
Taste dysfunction	54.2%
Sore throat	52.9%
Fever (>38°C)	45.4%

RNA = ribonucleic acid; C = Celsius.

Guan WJ, et al. *N Engl J Med.* 2020;382:1708-1720. Rothan HA, et al. *J Autoimmun.* 2020;109:102433. Lechien JR, et al. *J Intern Med.* 2020;288:335-344. Wang WW, et al. *J Med Virol.* 2020;92:441-447.

COVID-19 Disease Severity


A large study of 44,672 confirmed COVID-19 cases identified by the Chinese Centers for Disease Control and Prevention found that 81% of cases were mild-to-moderate, 13.8% were severe, and 6.1% were critical

	Disease Characteristics—NIH
Mild illness	Various symptoms (eg, fever, cough, sore throat, headache, malaise, muscle pain) without shortness of breath, dyspnea, or abnormal chest imaging
Moderate illness	SpO ₂ ≥94% on room air and lower respiratory disease evidenced by clinical assessment or imaging
Severe illness	SpO ₂ <94% on room air, PaO ₂ /FiO ₂ <300, respiratory rate >30 breaths/min, or lung infiltrates >50%
Critical illness	Respiratory failure, septic shock, and/or multiorgan dysfunction

SpO₂ = oxygen saturation; PaO₂ = arterial partial pressure of oxygen; FiO₂ = fraction of inspired oxygen; NIH = National Institutes of Health.

Wu Z, McGoogan JM. *JAMA.* 2020;323:1239-1242. NIH. COVID-19 treatment guidelines (<https://files.covid19treatmentguidelines.nih.gov/guidelines/covid19treatmentguidelines.pdf>). Accessed 12/2/2020.

Phases of COVID-19



ARDS = acute respiratory distress syndrome; CRP = C-reactive protein; IL-6 = interleukin 6; LDH = lactate dehydrogenase; NT-proBNP = N-terminal pro B-type natriuretic peptide; SIRS = systemic inflammatory response syndrome.

Adapted from Siddiqi HK, Mehra MR. *J Heart Lung Transplant.* 2020;39:405-407.

Association Between Pre-existing Characteristics and COVID-19 Survival

- Prospective cohort study of 20,133 patients in UK hospitalized with COVID-19
- Increasing age, male sex, and chronic comorbidities, including obesity, were identified as independent risk factors for mortality

		HR (95% CI)		P- value
Age on admission (years)	<50			
	50–59	2.63	(2.06–3.35)	<.001
	60–69	4.99	(3.99–6.25)	<.001
	70–79	8.51	(6.85–10.57)	<.001
	≥80	11.09	(8.93–13.77)	<.001
Sex at birth	Female	0.81	(0.75–0.86)	<.001
Chronic cardiac disease	Yes	1.16	(1.08–1.24)	<.001
Chronic pulmonary disease	Yes	1.17	(1.09–1.27)	<.001
Chronic kidney disease	Yes	1.28	(1.18–1.39)	<.001
Diabetes	Yes	1.06	(0.99–1.14)	.087
Obesity	Yes	1.33	(1.19–1.49)	<.001
Chronic neurological disorder	Yes	1.17	(1.06–1.29)	.001
Dementia	Yes	1.40	(1.28–1.52)	<.001
Malignancy	Yes	1.13	(1.02–1.24)	.017
Moderate/severe liver disease	Yes	1.51	(1.21–1.88)	<.001

UK = United Kingdom; HR = hazard ratio; CI = confidence interval.

Docherty AB, et al. *BMJ.* 2020;369:m1985.

Risk Factors for Severe Disease

Case series of 5700 hospitalized patients in NYC, Long Island, and Westchester County, NY found:

- Median number of total comorbidities at admission: 4 (IQR: 2–8)
- 88% of patients had more than one comorbidity
- Most common comorbidities were hypertension (56.6%), obesity (41.7%), and diabetes (33.8%)

Risk Factors for Severe COVID-19

- Older age
- Chronic obstructive pulmonary disease
- Cardiovascular disease (eg, heart failure, coronary artery disease, or cardiomyopathy)
- Type 2 diabetes mellitus
- Obesity (body-mass index >30)
- Sickle cell disease
- Chronic kidney disease
- Immunocompromised state from solid-organ transplantation
- Cancer

IQR = interquartile range.

Richardson S, et al. *JAMA*. 2020;323:2052-2059. NIH. COVID-19 treatment guidelines (<https://files.covid19treatmentguidelines.nih.gov/guidelines/covid19treatmentguidelines.pdf>). Accessed 12/2/2020.

Antibody Therapies for the Management of COVID-19

Antibody Therapy in Mild-to-Moderate COVID-19

	Asymptomatic or presymptomatic	Mild illness	Moderate illness	Severe illness	Critical illness
Features	Positive SARS-CoV-2 test; no symptoms	Mild symptoms (eg, fever, cough, or change in taste or smell); no dyspnea	Clinical or radiographic evidence of lower respiratory tract disease; oxygen saturation $\geq 94\%$	Oxygen saturation $< 94\%$; respiratory rate ≥ 30 breaths/min; lung infiltrates $> 50\%$	Respiratory failure, shock, and multiorgan dysfunction or failure
Testing	Screening testing; if patient has known exposure, diagnostic testing	Diagnostic testing	Diagnostic testing	Diagnostic testing	Diagnostic testing
Isolation	Yes	Yes	Yes	Yes	Yes
Proposed disease pathogenesis					
Potential treatment		Antiviral therapy	Antibody therapy	Anti-inflammatory therapy	
Management considerations	Monitoring for symptoms	Clinical monitoring and supportive care	Clinical monitoring; if patient hospitalized and at high risk for deterioration, possibly remdesivir	Hospitalization, oxygen therapy, and specific therapy (remdesivir, dexamethasone)	Critical care and specific therapy (dexamethasone, possibly remdesivir)

According to the Centers for Disease Control and Prevention (CDC), diagnostic testing for SARS-CoV-2 is intended to identify current infection in individuals and is performed when a person has signs or symptoms consistent with Covid-19 or when a person is asymptomatic but has recent known or suspected exposure to SARS-CoV-2. Screening testing for SARS-CoV-2 is intended to identify infected persons who are asymptomatic and without known or suspected exposure to SARS-CoV-2. Screening testing is performed to identify persons who may be contagious so that measures can be taken to prevent further transmission.


Adapted from Gandhi RT, et al. *N Engl J Med.* 2020;383:1757-1766

Convalescent Plasma in Severe COVID: Not Highly Effective

Convalescent Plasma Compared with Placebo

- Convalescent plasma had no impact on survival at 30 days in severe COVID-19 in 333 hospitalized patients

- >98% on convalescent plasma and 95% on placebo with oxygen saturation $< 93\%$ on room air at baseline
- 29% on convalescent plasma and 24% on placebo in ICU at baseline
- 92% on convalescent plasma and 96% on placebo given steroids during trial




ICU = intensive care unit.

Simonovich VA, et al. *N Engl J Med.* 2020;Nov 24: Epub ahead of print.

Effect of Anti-SARS-CoV-2 Antibody Level on 30-Day Mortality

- Death within 30 days after plasma transfusion in 3 titer groups
 - High-titer: 22.3% (115/515)
 - Medium-titer: 27.4% (549/2006)
 - Low-titer: 29.6% (166/561)
- Significantly lower risk of death within 30 days among patients who had not received mechanical ventilation before transfusion in high-titer group compared with low-titer group (RR = 0.66; 95% CI, 0.48–0.91)

High vs Low Antibody Levels



RR = relative risk; yr = year(s).

Joyner MJ, et al. *N Engl J Med*. 2021;Jan 13: Epub ahead of print.

Emergency Use Authorization (EUA) for Convalescent Plasma

- EUA issued for **high-titer** convalescent plasma
- Authorized for the treatment of hospitalized patients with COVID-19 early in the disease course and for hospitalized patients with impaired humoral immunity
- Early disease generally means prior to respiratory failure requiring intubation and mechanical ventilation

US Food and Drug Administration (FDA). Convalescent plasma fact sheet (www.fda.gov/media/141478/download). Accessed 2/25/2021.

Monoclonal Antibody Therapies

BLAZE-1: Phase 2 Trial of Bamlanivimab (LY-CoV555)

- Interim results from phase 2 trial of bamlanivimab in patients with mild-to-moderate COVID-19
- Risk factors for severe COVID-19 in 70% of bamlanivimab and 66% of placebo patients at baseline

Inclusion criteria:

- ≥ 18 years of age
- Not hospitalized
- Sample collection for 1st positive SARS-CoV-2 viral infection determination ≤ 3 days prior to start of infusion
- ≥ 1 mild or moderate symptom of COVID-19 (fever, cough, sore throat, malaise, headache, muscle pain, gastrointestinal symptoms, or shortness of breath with exertion)

R
N = 452

LY-CoV555 700 mg
monotherapy
(n = 101)

LY-CoV555 2800 mg
monotherapy
(n = 107)

LY-CoV555 7000 mg
monotherapy
(n = 101)

Placebo
(n = 143)

Interim analysis


- Positive SARS-CoV-2 test ≤ 3 days before infusion
- Mild or moderate COVID-19 symptoms
- Primary endpoint: change from baseline to day 11 (± 4 days) in SARS CoV-2 viral load
- Secondary endpoints include safety, symptom severity, hospitalization, and time points for viral clearance

BLAZE-1 Interim Results

Treatment	Patients Hospitalized/ Total No.	Incidence of Hospitalization (%)
Placebo	9/143	6.3
Bamlanivimab 700 mg	1/101	1.0
Bamlanivimab 2800 mg	2/107	1.9
Bamlanivimab 7000 mg	2/101	2.0
Bamlanivimab pooled doses	5/309	1.6

- In subjects ≥ 65 years and/or with a BMI ≥ 35 , day 29 hospitalization was 4% in treated patients and 15% in those receiving placebo

Symptom score from day 2 to day 11



- Symptom scores ranged from 0 to 24 and included eight domains, each of which was graded on a scale of 0 (no symptoms) to 3 (severe symptoms)


BMI = body-mass index.

Chen P, et al. *N Engl J Med.* 2020;Oct 28: Epub ahead of print.


BLAZE-1: Viral Loads Over Time

- Correlation between high viral load and hospitalization
- At day 7, the frequency of hospitalization was 12% (7 of 56 patients) among those who had a Ct value of less than 27.5, as compared with a frequency of 0.9% (3 of 340 patients) among those with a lower viral load.

Viral load in all patients



Viral load on day 7 in each trial group



Ct = PCR (polymerase-chain reaction) cycle threshold (higher viral load = lower Ct value).

Chen P, et al. *N Engl J Med.* 2020;Oct 28: Epub ahead of print.

BLAZE-1: Bamlanivimab Safety

- No serious AEs reported with bamlanivimab use


Adverse Event	LY-CoV555 (N=309)				Placebo (n = 143)
	700 mg (n = 101)	2800 mg (n = 107)	7000 mg (n = 101)	Pooled Doses (n = 309)	
Serious adverse event	0	0	0	0	1 (0.7)
Adverse events					
Any	24 (23.8)	23 (21.5)	22 (21.8)	69 (22.3)	35 (24.5)
Mild	16 (15.8)	18 (16.8)	10 (9.9)	44 (14.2)	18 (12.6)
Moderate	7 (6.9)	3 (2.8)	8 (7.9)	18 (5.8)	16 (11.2)
Severe	0	2 (1.9)	3 (3.0)	5 (1.6)	1 (0.7)
Missing data	1 (1.0)	0	1 (1.0)	2 (0.6)	0

- Infusion-related reactions were reported in 2.3% of patients receiving bamlanivimab and 1.4% of patients in the placebo group
 - Most reactions were mild and occurred during the infusion

Chen P, et al. *N Engl J Med.* 2020;Oct 28: Epub ahead of print.

ACTIV-3 Trial: Bamlanivimab in Hospitalized Patients

- Hospitalized patients were randomized to receive bamlanivimab or placebo in addition to high-quality supportive care, including remdesivir and, when indicated, supplemental oxygen and glucocorticoids




- Trial was paused when bamlanivimab was not shown to improve outcomes in hospitalized patients with COVID-19 who did not have end-organ failure

ACTIV-3/TICO LY-CoV555 Study Group. *N Engl J Med.* 2020;Dec 22: Epub ahead of print.

Emergence of SARS-CoV-2 Variants


- Several SARS-CoV-2 variants with enhanced transmissibility have emerged
 - B.1.1.7 contains 8 spike mutations and emerged in the UK
 - B.1.351 from South Africa has 9 spike mutations
- Activity against the B.1.351 variant is:
 - Reduced with casirivimab
 - Absent with bamlanivimab



WA1 = wild-type strain; UKΔ8 = pseudovirus with 8 B.1.1.7 mutations; SAΔ9 = pseudovirus with 9 B.1.351 mutations; IC50 = half maximal inhibitory concentration. Wang P, et al. *Nature*. 2021;Epub ahead of print.

Mechanism of Action of mAb Therapies Against SARS-CoV-2


- Neutralizing monoclonal antibodies against SARS-CoV-2 bind to the receptor-binding domain (RBD) of the spike protein and prevent host-cell entry
- Dual monoclonal antibody cocktail contains 2 potent antibodies that simultaneously and noncompetitively bind to different regions of the RBD
 - Use of 2 individual antibodies prevents generation of escape mutants and therapy failure



Hansen J, et al. *Science*. 2020;369:1010-1014.

Antibody Escape Mutations in Circulating SARS-CoV-2

- Many variants that can escape a single monoclonal antibody are currently in circulation
- Very few variants are capable of escaping dual monoclonal-antibody therapies



GISAID = Global Initiative on Sharing Avian Influenza Data.

Starr TN, et al. *Science*. 2021;371:850-854.

Bamlanivimab Plus Etesevimab

- Etesevimab is a neutralizing monoclonal antibody that binds to a different epitope on the spike protein than bamlanivimab
- 577 nonhospitalized patients with mild-to-moderate COVID-19 were randomized to bamlanivimab (700 mg, 2800 mg, or 7000 mg), combination therapy (bamlanivimab 2800 mg + etesevimab 2800 mg), or placebo

	Bamlanivimab 700 mg	Bamlanivimab 2800 mg	Bamlanivimab 7000 mg	Bamlanivimab 2800 mg + Etesevimab 2800 mg	Placebo
Change in log viral load from baseline to day 11	-3.72 <i>P</i> = 0.69	-4.08 <i>P</i> = 0.21	-3.49 <i>P</i> = 0.16	-4.37 <i>P</i> = 0.01	-3.80
COVID-19-related hospitalizations or ED visits	1.0%	1.9%	2.0%	0.9%	5.8%

ED = emergency department.

Gottlieb RL, et al. *JAMA*. 2021;Jan 21: Epub ahead of print.

Casirivimab and Imdevimab (REGN-COV2)

Ongoing phase 1–3 trial of casirivimab and imdevimab in nonhospitalized adults with mild-to-moderate COVID-19

Inclusion criteria:

- ≥18 years
- ≥1 symptom of COVID-19
- Positive SARS-CoV-2 test <72 hours prior to randomization
- Symptoms consistent with COVID-19 with onset <7 days before randomization
- No hospitalization due to COVID-19

R

2.4 g casirivimab and imdevimab
(1.2 g each)

8.0 g casirivimab and imdevimab
(4.0 g each)

Placebo

Casirivimab and imdevimab EUA. (www.fda.gov/media/143892/download). Accessed 1/7/2021.

Casirivimab and Imdevimab: Interim Results

Interim analysis of 275 nonhospitalized patients with mild-to-moderate COVID-19

At Least 1 COVID-19-Related Medical Visit Within 29 Days		
Treatment	Events/Total Patients	Incidence
All patients		
Placebo	6/93	6%
Casirivimab and imdevimab 2.4 g	3/92	3%
Casirivimab and imdevimab 8.0 g	3/90	3%
All doses casirivimab and imdevimab	6/182	3%
Seronegative patients		
Placebo	5/33	15%
Casirivimab and imdevimab 2.4 g	2/41	5%
Casirivimab and imdevimab 8.0 g	3/39	8%
All doses casirivimab and imdevimab	5/80	6%

Weinreich DM, et al. *N Engl J Med.* 2020;Dec 17: Epub ahead of print.

Casirivimab/Imdevimab: Efficacy by Baseline Viral Load

Casirivimab/imdevimab (REGN-COV2) provided greater reduction in viral load in those patients with higher viral load at baseline





Viral load over time according to baseline viral-load category

	>10 ⁴ copies/mL	
	Difference in Change from Baseline, Day 7	
	TWA LS mean	Mean
2.4 g vs PBO	-0.36	-0.64
8.0 g vs PBO	-0.59	-0.90

	>10 ⁵ copies/mL	
	Difference in Change from Baseline, Day 7	
	TWA LS mean	Mean
2.4 g vs PBO	-0.59	-0.83
8.0 g vs PBO	-0.75	-1.12

	>10 ⁶ copies/mL	
	Difference in Change from Baseline, Day 7	
	TWA LS mean	Mean
2.4 g vs PBO	-0.81	-1.46
8.0 g vs PBO	-1.14	-1.54

	>10 ⁷ copies/mL	
	Difference in Change from Baseline, Day 7	
	TWA LS mean	Mean
2.4 g vs PBO	-1.03	-1.84
8.0 g vs PBO	-1.32	-1.75



TWA = time-weighted average; LS = least-squares.

Weinreich DM, et al. *N Engl J Med.* 2020;Dec 17: Epub ahead of print.

Casirivimab/Imdevimab Safety

Event	REGN-COV2			Placebo (n = 93)
	2.4 g (n = 88)	8.0 g (n = 88)	Combined (n = 176)	
	Number of patients (%)			
Any serious adverse event	1 (1)	0	1 (1)	2 (2)
Any adverse event of special interest* (Grade 2 or higher hypersensitivity or infusion-related reactions)	0	2 (2)	2 (1)	2 (2)
Any serious adverse event of special interest*	0	0	0	0
Grade ≥2 infusion-related reaction within 4 days	0	2 (2)	2 (1)	1 (1)
Grade ≥2 hypersensitivity reaction within 29 days	0	1 (1)	1 (1)	2 (2)
Adverse events that occurred or worsened during the observation period†				
Grade 3 or 4 event	1 (1)	0	1 (1)	1 (1)
Event that led to death	0	0	0	0
Event that led to withdrawal from the trial	0	0	0	0
Event that led to infusion interruption*	0	1 (1)	1 (1)	1 (1)

*Events were grade 2 or higher hypersensitivity reactions or infusion-related reactions.

†Events listed here were not present at baseline or were an exacerbation of a preexisting condition that occurred during the observation period, which is defined as the time from administration of REGN-COV2 or placebo to the last study visit.

Weinreich DM, et al. *N Engl J Med.* 2020;Dec 17: Epub ahead of print.

mAb Therapies With Emergency Use Authorization

These therapies must be given as soon as possible
and within 10 days of symptom onset

**Bamlanivimab 700 mg
AND
Etesevimab 1400 mg**

Administer together as
single IV infusion over
minimum of 21–60
minutes

**Casirivimab 1200 mg
AND
Imdevimab 1200 mg**

Must be administered
together as a single IV
infusion over minimum
of 60 minutes

**Monotherapy not
recommended due to
resistance of viral variants**

Bamlanivimab 700 mg

Administer as a single
IV infusion over
minimum of 16–60
minutes

IV = Intravenous.

Bamlanivimab EUA. (<http://pi.lilly.com/eua/bamlanivimab-eua-factsheet-hcp.pdf>). Bamlanivimab and etesevimab EUA. (www.fda.gov/media/145802/download). Casirivimab and imdevimab EUA. (www.fda.gov/media/143892/download). US Health and Human Services (HHS). (www.phe.gov/emergency/events/COVID19/investigation-MCM/Bamlanivimab/Pages/default.aspx)

Emergency Use Authorization of COVID-19 mAb Therapy

- EUA for the treatment of mild-to-moderate COVID-19 in patients:
 - Who are at least 12 years of age and weigh at least 40 kg
 - Have positive results of direct SARS-CoV-2 viral testing
 - Who are at high risk of progressing to severe COVID-19 or hospitalization
- No benefit in patients hospitalized due to COVID-19
- These therapies may be associated with worse clinical outcomes in hospitalized COVID-19 patients requiring high-flow oxygen or mechanical ventilation

Casirivimab and imdevimab EUA. (www.fda.gov/media/143892/download). Bamlanivimab EUA. (<http://pi.lilly.com/eua/bamlanivimab-eua-factsheet-hcp.pdf>). URLs accessed 12/2/2020.

Identifying High-Risk Candidates for mAb Therapy

High risk is defined as a patient who meets ≥ 1 of the following criteria

Patients of any age with:

- BMI ≥ 35
- Chronic kidney disease
- Diabetes
- Immunosuppressive disease
- Current immunosuppressive therapy

Patients aged ≥ 65 years

Patients ≥ 55 years of age with:

- Cardiovascular disease, OR
- Hypertension, OR
- Chronic obstructive pulmonary disease/
other chronic respiratory condition

Patients aged 12–17 years with:


- BMI >85th percentile for age and gender
- Sickle cell disease
- Congenital or acquired heart disease
- Neurodevelopmental disorders (eg, cerebral palsy)
- Asthma, reactive airway, or other chronic respiratory disease that requires daily medication for control
- A medical-related technological dependence (eg, tracheostomy, gastrostomy, positive-pressure ventilation not related to COVID-19)

Casirivimab and imdevimab EUA. (www.fda.gov/media/143892/download). Bamlanivimab EUA. (<http://pi.lilly.com/eua/bamlanivimab-eua-factsheet-hcp.pdf>). URLs accessed 12/2/2020.

Bamlanivimab in Nursing-Home Setting

- 966 participants, including 266 nursing-home residents considered at high-risk for severe COVID-19, were administered a single-dose of bamlanivimab or placebo if a case of SARS-CoV-2 was confirmed in nursing home
- Compared with placebo, bamlanivimab was associated with:
 - Significantly lower proportion of residents with mild or worse COVID-19 by day 57 (OR = 0.20; 95% CI, 0.08–0.49; $P < .001$)
 - Significant reductions in incident SARS-CoV-2 infection by day 29 (OR = 0.23; CI, 0.11–0.48; $P < .001$)
- 5 COVID-19-related deaths (all in placebo group)

Time since treatment to development of mild or worse COVID-19 in residents



OR = odds ratio.

Cohen M, et al. CROI 2021: abstract 1211B. Lilly BLAZE-2 press release. 1/21/2121. (<https://investor.lilly.com/node/44291/pdf>). Accessed 3/25/2021.

Top-line Results on mAb Therapies

- **BLAZE-1: Bamlanivimab plus etesivimab**
 - Phase 3 trial of 769 high-risk, recently diagnosed COVID-19 patients showed that therapy with bamlanivimab and etesevimab reduced hospitalizations and deaths by **87%** ($P= .0001$)
- **Casirivimab and imdevimab for COVID-19 treatment**
 - 70% reduction in risk of hospitalization or death in 4567 high-risk, non-hospitalized COVID-19 patients
- **Casirivimab and imdevimab for COVID-19 prevention**
 - Interim analysis found 100% prevention of symptomatic infection and 50% reduction in rate of COVID-19 infection in a phase 3 trial of 400 individuals with household exposure to COVID-19

Lilly press release. 3/10/2021. (<https://investor.lilly.com/news-releases/news-release-details/lillys-bamlanivimab-and-etesevimab-together-reduced>). Regeneron press release. 1/26/21. (<https://newsroom.regeneron.com/news-releases/news-release-details/regeneron-reports-positive-interim-data-regen-covtm-antibody>). Regeneron press release. 3/23/21. (<https://investor.regeneron.com/news-releases/news-release-details/phase-3-trial-shows-regen-covtm-casirivimab-imdevimab-antibody>). URLs accessed 3/25/2021.

COVID-19 Antibody Treatment Resource Guide National Infusion Center Association

- Infusion center locator
- Resources for providers
 - Bamlanivimab playbook
 - Casirivimab + imdevimab guidebook
- Patient education resources
- Treatment indication checklist
- Plus, other resources

The screenshot shows the title page of the "COVID-19 Antibody Treatment Resource Guide" from the National Infusion Center Association (NICA). The page includes the NICA logo and a brief introduction stating that the resources are developed to support prescribers, infusion providers, and patients. It lists several key resources: the NICA COVID-19 Locator, a section on "Locating Sites of Care" with instructions for prescribers and patients, and the HHS Protect Public Data Hub for Therapeutics Distribution Locations. The locator section provides detailed steps for finding infusion sites, including using a search tool and providing specific information for prescribers and patients.

National Infusion Center Association (https://infusioncenter.org/infusion_resources/covid-19-antibody-treatment-resource-center/). Accessed 1/18/2021.

Management of Hospitalized Patients with COVID-19

IDSA: Recommended Treatment Options for Hospitalized Patients

Treatment	Guidance
Remdesivir	<ul style="list-style-type: none"> • Recommended for hospitalized patients with severe COVID-19 • Most benefit seen in those with severe COVID-19 on supplemental oxygen rather than patients on mechanical ventilation or ECMO • 5 days of treatment recommended for patients on supplemental oxygen • 10 days of treatment recommended for patients on mechanical ventilation or ECMO
Glucocorticoids	<ul style="list-style-type: none"> • Recommended for hospitalized patients with severe COVID-19 • Dexamethasone 6 mg IV or PO for 10 days or equivalent • Not recommended for hospitalized patients without hypoxemia ($SpO_2 >94\%$) requiring supplemental oxygen
Baricitinib plus remdesivir	<ul style="list-style-type: none"> • Baricitinib plus remdesivir recommended over remdesivir alone in hospitalized patients with severe COVID-19 who cannot receive corticosteroids because of a contraindication
Tocilizumab	<ul style="list-style-type: none"> • Recommended in addition to standard of care in hospitalized patients with progressive severe or critical COVID-19 who have elevated markers of systemic inflammation

IDSA = Infectious Diseases Society of America; ECMO = extracorporeal membrane oxygenation; PO = by mouth.


Bhimraj A, et al. IDSA Guidelines. V3.9.0. (www.idsociety.org/practice-guideline/covid-19-guideline-treatment-and-management/).

Adaptive COVID-19 Treatment Trial (NIAID ACTT-1): Trial Design

- Multicenter, adaptive, randomized, double-blind, placebo-controlled phase 3 trial

Inclusion criteria (N = 1062)

- Adult patients ≥ 18 years of age
- Hospitalized with symptoms of COVID-19/SARS-CoV-2 infection and ≥ 1 of following:
 - Radiographic infiltrates by imaging
 - $SpO_2 \leq 94\%$ on room air
 - Requiring supplemental oxygen
 - Requiring mechanical ventilation



- Primary endpoint: time to recovery by day 29 according to 8-point ordinal scale
- Secondary endpoints: treatment-related improvements in ordinal scale at day 15

QD = each day.

Beigel JH, et al. *N Engl J Med.* 2020;383:1813-1826.

COVID-19 Clinical Status Ordinal Scale

Clinical Status Ordinal Scale	Clinical Status Description for Assessment
1	Not hospitalized, no limitations on activities
2	Not hospitalized, limitation on activities, and/or requiring home oxygen
3	Hospitalized, not requiring supplemental oxygen, and no longer requires ongoing medical care (if hospitalization extended for infection-control purposes)
4	Hospitalized, not requiring supplemental oxygen; requiring ongoing medical care (COVID-19 related or otherwise)
5	Hospitalized, requiring supplemental oxygen
6	Hospitalized, on noninvasive ventilation or high-flow oxygen devices
7	Hospitalized, on invasive mechanical ventilation or ECMO
8	Death

Beigel JH, et al. *N Engl J Med.* 2020;383:1813-1826.

Remdesivir: NIAID ACTT Clinical Trial

- 1062 patients in 68 sites randomized 1:1 to remdesivir or placebo
- Independent data safety monitoring board found that remdesivir shortened time to recovery compared with placebo

	Remdesivir	Placebo	P-value
Time to recovery	10 days	15 days	P < .001
Mortality	6.7% day 15 11.4% day 29	11.9% day 15 15.2% day 29	P = .07 (day 29)




An ICU bed becomes available
5 days earlier
Benefit is in early disease



~30% reduction in mortality
Not statistically significant

NIAID = National Institute of Allergy and Infectious Diseases.
Beigel JH et al. *N Engl J Med.* 2020;383:1813-1826 plus supplement.


Remdesivir: NIAID ACTT Clinical Trial



Beigel JH, et al. *N Engl J Med.* 2020;383:1813-1826.

WHO SOLIDARITY Trial: Antiviral Drugs in Hospitalized Patients

Open-label, randomized phase 3 trial conducted in 405 hospitals in 30 countries




- **Primary endpoint**
28-day in-hospital mortality
- **Secondary endpoints**
 - Initiation of ventilation
 - Duration of hospitalization

BID = twice daily; SQ = subcutaneous.

Pan H, et al; WHO (World Health Organization) Solidarity Trial Consortium. *N Engl J Med.* 2021;384:497-511.

WHO SOLIDARITY Trial: Results

Remdesivir, hydroxychloroquine, lopinavir, and interferon regimens had little to no effect on overall mortality, initiation of ventilation, or duration of hospital stay in hospitalized patients with COVID-19



Pan H, et al; WHO Solidarity Trial Consortium. *N Engl J Med.* 2021;384:497-511.

RECOVERY Trial Design

- Eligible patients (hospitalized with clinically suspected or laboratory-confirmed SARS-CoV-2 infection) were randomized to:

No additional treatment

Dexamethasone

Hydroxychloroquine

Lopinavir/ritonavir

Azithromycin


- Primary endpoint: 28-day mortality
- Patients with progressive disease (hypoxia and an inflammatory state) may undergo second randomization to no additional treatment or **tocilizumab**
- Current RECOVERY trials are investigating **baricitinib**, **casirivimab/imdevimab**, **aspirin**, **dexamethasone** (in children), and **colchicine**

Randomized Evaluation of COVID-19 Therapy—RECOVERY (www.recoverytrial.net/files/recovery-protocol-v7-0-2020-06-18.pdf). Accessed 2/12/2021.

RECOVERY Trial

Mortality With Dexamethasone + UC vs UC Alone

2104 patients randomized to dexamethasone 6 mg QD for up to 10 days; 4321 patients received UC alone




UC = usual care.

Horby P, et al; RECOVERY Collaborative Group. *N Engl J Med.* 2020;Jul 17:Epub ahead of print.

RECOVERY Trial


Mortality in Patients On Oxygen or Mechanical Ventilation ± Dexamethasone

Oxygen only (n = 3883)



No. at risk		0	7	14	21	28
Dexamethasone/UC	1279	1135	1036	1006	981	
UC	2604	2195	2018	1950	1916	

Invasive mechanical ventilation (n = 1007)



No. at Risk		0	7	14	21	28
Dexamethasone/UC	324	290	248	232	228	
UC	683	572	481	424	400	

Horby P, et al; RECOVERY Collaborative Group. *N Engl J Med.* 2020;Jul 17:Epub ahead of print.

EUAs for Hospitalized Patients

ACCT-2: Baricitinib Plus Remdesivir in Hospitalized Patients

Inclusion criteria:

- ≥18 years
- Hospitalized with COVID-19
- ≥1 of the following criteria:
 - Radiographic infiltrates by imaging
 - SpO₂ ≤94% on room air
 - Supplemental oxygen, mechanical ventilation, or ECMO

N =
1033

Baricitinib (≤14 days) plus
remdesivir (≤10 days)
(n = 515)

Placebo plus remdesivir (≤10 days)
(n = 518)

- Primary outcome: time to recovery
- Key secondary outcome: clinical status at day 15

Kalil AC, et al. *N Engl J Med.* 2021;384:795-807.


ACTT-2: Ordinal Scale

Ordinal Scale Used for Outcome Measures	
Recovered	
1	Not hospitalized, no limitations on activities
2	Not hospitalized, limitation on activities and/or requiring home oxygen
3	Hospitalized, not requiring supplemental oxygen—no longer requiring ongoing medical care
Population enrolled	
4	Hospitalized, not requiring supplemental oxygen—requiring ongoing medical care
5	Hospitalized, requiring supplemental oxygen
6	Hospitalized, on non-invasive ventilation or high-flow oxygen devices
7	Hospitalized, on mechanical ventilation or ECMO
8	Death

Kalil AC, et al. *N Engl J Med.* 2021;384:795-807.

Baricitinib Plus Remdesivir: Recovery Time

- Recovery time was reduced with baricitinib vs placebo (7 days vs 8 days; rate ratio for recovery = 1.16; 95% CI, 1.01–1.32; $P = .03$)



- Time to recovery was significantly lower with baricitinib in patients receiving high-flow oxygen or noninvasive ventilation at enrollment (10 days vs 18 days; rate ratio for recovery = 1.51)

RDV = remdesivir.

Kalil AC, et al. *N Engl J Med*. 2021;384:795-807.

Baricitinib Plus Remdesivir: Results

- Baricitinib was associated with 30% higher odds of improvement in clinical status at day 15 (OR = 1.3)
- 28-day mortality was 5.1% in the combination group and 7.8% in the control group (HR for death = 0.65)

Overall Outcomes		
Outcomes	Baricitinib + RDV (n = 515)	Placebo + RDV (n = 518)
Recovery		
No. of recoveries	433	406
Median time to recovery (95% CI), days	7 (6–8)	8 (7–9)
Rate ratio (95% CI)	1.16 (1.01–1.32), $P = .03$	
Mortality over first 14 days		
No. of deaths by day 14	8	15
Kaplan-Meier estimate of mortality by day 14, % (95% CI)	1.6 (0.8–3.2)	3.0 (1.8–5.0)
HR (95% CI) for data through day 14	0.54 (0.23–1.28)	
Mortality over entire trial period		
No. of deaths by day 28	24	37
Kaplan-Meier estimate of mortality by day 28, % (95% CI)	5.1 (3.5–7.6)	7.8 (5.7–10.6)
HR (95% CI)	0.65 (0.39–1.09)	

Kalil AC, et al. *N Engl J Med*. 2021;384:795-807.

ACTT-2: Adverse Events

Treatment-Emergent Adverse Events in ACTT-2		
	Baricitinib + RDV (n = 508) No. (%)	Placebo + RDV (n = 509) No. (%)
Grade 3 or 4 AEs	207 (40.7)	238 (46.8)
Hyperglycemia	25 (4.9)	40 (7.9)
Anemia	25 (4.9)	33 (6.5)
Decreased lymphocyte count	24 (4.7)	35 (6.9)
Acute kidney injury	20 (3.9)	36 (7.1)
Venous thromboembolism	21 (4.1)	16 (3.1)

AE = adverse event.

Kalil AC, et al. *N Engl J Med.* 2021;384:795-807.

Emergency Use Authorization for Baricitinib

- Baricitinib plus remdesivir was authorized for emergency use in hospitalized adults and pediatric patients ≥ 2 years of age requiring supplemental oxygen, invasive mechanical ventilation, or ECMO with suspected or confirmed COVID-19
- Recommended dosage:
 - Patients ≥ 9 years of age: 4 mg baricitinib once daily
 - Patients 2 to 9 years of age: 2 mg baricitinib once daily
- Recommended treatment duration is 14 days or until hospital discharge, whichever comes first
- Evaluate baseline eGFR, liver enzymes, and complete blood count to determine treatment suitability and dose

Baricitinib EUA. (www.fda.gov/media/143823/download). Accessed 1/20/2021.

Case Study

Early Intubation

History of Present Illness

- BW is 65-year old black male who presents with a chief complaint of profound fatigue beginning 3 days prior to admission
 - Patient was unable to get out of bed and had body aches, anorexia, and chills
 - Developed mild shortness of breath, rhinorrhea and dry cough within last 24 hours
 - No headaches, sore throat, postnasal drip, ear pain, chest pain, palpitation, or loss of taste/smell
- Primary care physician reported 102.7° fever and O₂ saturation of 89% on room air, and referred patient to the ED

Past Medical History and Medications

- Past medical history significant for:

- Hypertension
- Hyperlipidemia
- Chronic kidney disease
- Type 2 diabetes
- Gout
- Hyperparathyroidism
- Obstructive sleep apnea


Medication	Dosage
Allopurinol	10 mg daily
Amlodipine	10 mg daily
Carvedilol	6.25 mg BID with meals
Chlorthalidone	25 mg daily
Colchicine	0.6 mg PRN for gout
Ergocalciferol	50,000 units weekly
Glipizide	5 mg ER daily
Metformin	500 mg XR daily
Minoxidil	5 mg BID

Is BW a candidate for monoclonal antibody therapy?

Hospital Admission

- At the ED, patient was found to be hypoxic with SpO₂ of 80% on room air and he required 6 L NC with bilateral opacities on CXR
- He was admitted and moved to the ICU 2 days later after developing tachypnea and an altered mental status
 - 23 breaths per minute
 - Elevated inflammatory markers (CRP: 261 mg/L, ferritin: 999.6 ng/mL)
 - eGFR: 30 mL/min/1.73m²
- Due to persistent hypoxia (SpO₂ of 84% on 2 L NC), decision was made for early intubation

Imaging and Lab Results



- Imaging:
 - Widespread peribronchial opacities
 - Cardiomegaly
- Lab Results:
 - Sputum culture: negative
 - Strep pneumo Ag: negative
 - Flu A/B: negative
 - Blood cultures: no growth

How would you manage BW?

Summary of Hospital Course

- ARDS treated with a 10 day course of dexamethasone
- Patient developed Pseudomonas bacteremia in hospital and completed IV cefepime course
- Acute renal failure that developed initially required intermittent hemodialysis but eventually resolved
- Extubated after 14 days of ventilator support
- Discharged to home after 39 days of hospitalization

Case Study

Severe COVID-19 in a Pregnant Woman

History of Present Illness

- RD is a 33-year-old woman (G5P2) who is 28 weeks 2 days pregnant
- She presents with shortness of breath and fatigue with intermittent subjective fever, chills, diaphoresis, and headache that began 5 days ago
- Prior medical history significant for:
 - Type 2 diabetes (latest HbA1c 7.9%) managed with insulin glargine 120 units daily, insulin aspart 82/68/70, and metformin 1000 mg BID
 - Obesity (BMI 34)
 - Gestational hypertension
 - History of 2 prior Caesarian sections and 2 spontaneous abortions

Is RD a candidate for monoclonal antibody therapy?

Hospital Admission

- Patient was admitted but did not require supplemental oxygen
 - She tested positive for SARS-CoV-2
 - CXR was negative
 - Mild shortness of breath and mild transaminitis (ALT 75, AST 140) noted
- She was discharged 2 days later but was readmitted with worsening shortness of breath 3 days later
 - She was found to be hypoxic with O₂ saturation dropping to 80% on ambulation
 - She required 6-8L NC O₂ support
 - Denies sore throat, nasal congestion, loss of taste or smell, or GI symptoms
 - Review of systems were all negative

Lab and Imaging Results

- WBC: 6.0 (N85% L12%) w/ lymphopenia (700 cells/ μ L)
- Hb: 11.5
- Platelets: 252
- Na: 137, K: 4.7, Cl: 110, CO₂: 11 (ABG 16)
- Glucose: 247
- BUN: 6, Creatinine: 0.43 (eGFR >60)
- ALT: 48, AST: 62, ALP: 99, TB: 0.5
- CRP: 20.1 (normal: \leq 1 mg/dL)
- LDH: 606
- D-dimer: 580 (normal: <500 ng/mL)
- Ferritin: 266.8 (normal: 6.2-137 ng/mL)



Audience Question

How would you manage this patient?

Hospital Course

- RD required HFNC and BiBAP on the 1st hospital day (HD) and was ultimately intubated on her 4th HD
- Patient was treated with a 10-day course of remdesivir and received betamethasone on the 1st and 2nd HD due to concern for preterm labor
- Her course was complicated by multi-drug resistant *Acinetobacter baumannii* ventilator-associated pneumonia
 - Completed 7 days of ampicillin/sulbactam, meropenem, and inhaled colistin
- She subsequently improved
 - Extubated on 10th HD and discharged on 22nd HD

Summary

- Several neutralizing mAb therapies are authorized for treatment of mild-to-moderate COVID-19 in patients at high risk of progressing to severe COVID-19 or hospitalization
 - mAbs against SARS-CoV-2 reduced the risk of COVID-19-related hospitalization
 - These therapies may be associated with worse clinical outcomes in hospitalized COVID-19 patients requiring high-flow oxygen or mechanical ventilation
 - Therapy should be provided as soon as possible and within 10 days of symptoms onset
 - Due to the emergence of viral resistance, dual monoclonal antibody therapies—bamlanivimab plus etesevimab or casirivimab plus imdevimab—should be used
- Baricitinib plus remdesivir is authorized for emergency use in hospitalized adults and pediatric patients ≥ 2 years of age requiring supplemental oxygen, invasive mechanical ventilation, or ECMO with suspected or confirmed COVID-19
 - Recommended treatment duration is 14 days or until hospital discharge, whichever comes first
 - Baricitinib plus remdesivir associated with improvements in recovery time

Thank you!

A Light in the Darkness: New Virus-neutralizing Monoclonal Antibodies and Other Point-of-Care Therapies Recently Granted Emergency Use Authorizations for Patients with COVID-19

Resource	Address
National Infusion Center Association (NICA). COVID-19 Antibody Therapies Resource Center . Accessed February 23, 2021.	https://infusioncenter.org/infusion_resources/covid-19-antibody-treatment-resource-center/
Wiersinga WJ, et al. Pathophysiology, transmission, diagnosis, and treatment of coronavirus disease 2019 (COVID-19): A review . <i>JAMA</i> . 2020;324:782-793.	https://pubmed.ncbi.nlm.nih.gov/32648899/
Guan WJ, et al. Clinical characteristics of coronavirus disease 2019 in China . <i>N Engl J Med</i> . 2020;382:1708-1720.	https://pubmed.ncbi.nlm.nih.gov/32109013/
Rothan HA, et al. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak . <i>J Autoimmun</i> . 2020;109:102433.	https://pubmed.ncbi.nlm.nih.gov/32113704/
Lechien JR, et al. Clinical and epidemiological characteristics of 1420 European patients with mild-to-moderate coronavirus disease 2019 . <i>J Intern Med</i> . 2020;288:335-344.	https://pubmed.ncbi.nlm.nih.gov/32352202/
Wang W, et al. Updated understanding of the outbreak of 2019 novel coronavirus (2019-nCoV) in Wuhan, China . <i>J Med Virol</i> . 2020;92:441-447.	https://pubmed.ncbi.nlm.nih.gov/31994742/
Wu Z, et al. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: Summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention . <i>JAMA</i> . 2020;323:1239-1242.	https://jamanetwork.com/journals/jama/fullarticle/2762130
Richardson S, et al. Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York City area . <i>JAMA</i> . 2020.;323:2052-2059.	https://pubmed.ncbi.nlm.nih.gov/32320003/
Docherty AB, et al. Features of 20,133 UK patients in hospital with COVID-19 using the ISARIC WHO Clinical Characterisation Protocol: Prospective observational cohort study . <i>BMJ</i> . 2020;369:m1985.	https://www.bmj.com/content/369/bmj.m1985
Yuan X, et al. Changes of hematological and immunological parameters in COVID-19 patients . <i>Int J Hematol</i> . 2020;112:553-559.	https://pubmed.ncbi.nlm.nih.gov/32656638/
Bhimraj A, et al. Infectious Diseases Society of America. IDSA Guidelines on the Treatment and Management of Patients with COVID-19 . Version 4.0.0. Published 4/11/2020. Last updated 2/22/2021. Accessed February 23, 2021.	https://www.idsociety.org/COVID19guidelines

Alhazzani W, et al. Surviving Sepsis Campaign: Guidelines on the management of critically ill adults with coronavirus disease 2019 (COVID-19). <i>Intensive Care Med.</i> 2020;46:854-887.	https://pubmed.ncbi.nlm.nih.gov/32222812/
World Health Organization (WHO). Clinical management of COVID-19. Interim Guidance. Published May 27, 2020. Accessed February 23, 2021.	https://www.who.int/publications/i/item/clinical-management-of-covid-19
National Institutes of Health (NIH). COVID-19 Treatment Guidelines. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. Last updated February 23, 2021. Accessed February 23, 2021.	https://www.covid19treatmentguidelines.nih.gov/
Beigel JH, et al. Remdesivir for the treatment of COVID-19 – Final Report. <i>N Engl J Med.</i> 2020;383:1813-1826.	https://www.nejm.org/doi/full/10.1056/NEJMoa2007764
Spinner CD, et al. Effect of remdesivir vs standard care on clinical status at 11 days in patients with moderate COVID-19: A randomized clinical trial. <i>JAMA.</i> 2020;324:1048-1057.	https://jamanetwork.com/journals/jama/fullarticle/2769871
Horby P, et al. Dexamethasone in hospitalized patients with COVID-19 – preliminary report [published online ahead of print, 2020 Jul 17]. <i>N Engl J Med.</i> 2020; NEJMoa2021436.	https://pubmed.ncbi.nlm.nih.gov/32678530/
Siddiqi HK, et al. COVID-19 illness in native and immunosuppressed states: A clinical-therapeutic staging proposal. <i>J Heart Lung Transplant.</i> 2020;39:405-407.	https://www.jhltonline.org/article/S1053-2498(20)31473-X/fulltext
Li L, et al. Effect of convalescent plasma therapy on time to clinical improvement in patients with severe and life-threatening COVID-19: A randomized clinical trial. <i>JAMA.</i> 2020;324:460-470.	https://pubmed.ncbi.nlm.nih.gov/32492084/
Kalil AC, et al. Baricitinib plus remdesivir for hospitalized adults with COVID-19 [published online ahead of print, 2020 Dec 11]. <i>N Engl J Med.</i> 2020;NEJMoa2031994.	https://www.nejm.org/doi/full/10.1056/NEJMoa2031994
Baum A, et al. Antibody cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape seen with individual antibodies. <i>Science.</i> 2020;369:1014-1018.	https://pubmed.ncbi.nlm.nih.gov/32540904/
Gandhi RT, et al. Mild or moderate COVID-19. <i>N Engl J Med.</i> 2020;383:1757-1766.	https://pubmed.ncbi.nlm.nih.gov/32329974/
Docherty AB, et al. Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: Prospective observational cohort study. <i>BMJ.</i> 2020;369:m1985.	https://pubmed.ncbi.nlm.nih.gov/32444460/
Simonovich VA, et al. A Randomized trial of convalescent plasma in Covid-19 severe pneumonia. <i>N Engl J Med.</i> 2021;384:619-629.	https://pubmed.ncbi.nlm.nih.gov/33232588/

<p>US Food and Drug Administration (FDA). Fact Sheet for Health Care Providers. Emergency Use Authorization (EUA) of Baricitinib. Last updated November 19, 2020. Accessed February 23, 2021.</p>	<p>https://www.fda.gov/media/143823/download</p>
<p>US Food and Drug Administration (FDA). Fact Sheet for Health Care Providers. Emergency Use Authorization (EUA) of Bamlanivimab. Last updated February 9, 2021. Accessed February 23, 2021.</p>	<p>https://www.fda.gov/media/143603/download</p>
<p>US Food and Drug Administration (FDA). Fact Sheet for Health Care Providers. Emergency Use Authorization (EUA) of Casirivimab and Imdevimab. Last updated December 2020. Accessed February 23, 2021.</p>	<p>https://www.fda.gov/media/143892/download</p>
<p>Weinreich DM, et al. REGN-COV2, a neutralizing antibody cocktail, in outpatients with COVID-19. <i>N Engl J Med</i>. 2020;384:238-251.</p>	<p>https://www.nejm.org/doi/pdf/10.1056/NEJMoa2035002</p>
<p>Chen P, et al. SARS-CoV-2 neutralizing antibody LY-CoV555 in outpatients with Covid-19. <i>N Engl J Med</i>. 2020;384:229-237.</p>	<p>https://www.nejm.org/doi/full/10.1056/NEJMoa2029849</p>
<p>ACTIV-3/TICO LY-CoV555 Study Group, et al. A neutralizing monoclonal antibody for hospitalized patients with COVID-19 [published online ahead of print, 2020 Dec 22]. <i>N Engl J Med</i>. 2020; NEJMoa2033130.</p>	<p>https://www.nejm.org/doi/full/10.1056/NEJMoa2033130</p>
<p>Hansen J, et al. Studies in humanized mice and convalescent humans yield a SARS-CoV-2 antibody cocktail. <i>Science</i>. 2020;369:1010-1014.</p>	<p>https://science.sciencemag.org/content/369/6506/1010</p>
<p>Callaway E. The coronavirus is mutating – does it matter? <i>Nature</i>. 2020;585:174-177.</p>	<p>https://www.nature.com/articles/d41586-020-02544-6</p>