HM-52L: Updates for VA Clinicians in the Use of Bispecific Antibodies Across Later-Line Non-Hodgkin Lymphomas: Summarizing Current Efficacy and Safety Data

Toolkit

Introduction: Overview of B Cell Non-Hodgkin Lymphomas

Reference	URL
American Cancer Society (ACS). Key statistics for non-	https://www.cancer.org/cancer/types/non-
Hodgkin lymphoma. 2023.	hodgkin-lymphoma/about/key-statistics.html
Armitage JO, et al. New approach to classifying non- Hodgkin's lymphomas: Clinical features of the major	https://ascopubs.org/doi/10.1200/JCO.1998.16.8.
histologic subtypes. Non-Hodgkin's Lymphoma	
Classification Project. <i>J Clin Oncol</i> . 1998;16:2780-2795.	<u>2780</u>
Abramson JS, et al. Advances in the biology and therapy	https://ashpublications.org/blood/article/106/4/1
of diffuse large B-cell lymphoma: Moving toward a molecularly targeted approach. <i>Blood</i> . 2005;106:1164-	164/21561/Advances-in-the-biology-and-therapy-
1174.	<u>of-diffuse</u>
Baeuerle PA, et al. Bispecific T-cell engaging antibodies	https://aacrjournals.org/cancerres/article/69/12/4
for cancer therapy. Cancer Res. 2009 Jun 15;69(12):4941-	941/549633/Bispecific-T-Cell-Engaging-Antibodies-
4.	for-Cancer
Cerhan JR. Epidemiology of Follicular Lymphoma.	https://www.sciencedirect.com/science/article/ab
Hematol Oncol Clin North Am. 2020;34:631-646.	<u>s/pii/S0889858820300265</u>
Coiffier B, et al. CHOP chemotherapy plus rituximab	
compared with CHOP alone in elderly patients with	https://www.nejm.org/doi/full/10.1056/NEJMoa0
diffuse large-B-cell lymphoma. N Engl J Med.	<u>11795</u>
2002;346:235-242.	
Coiffier B, et al. Long-term outcome of patients in the	
LNH-98.5 trial, the first randomized study comparing	https://ashpublications.org/blood/article/116/12/
rituximab-CHOP to standard CHOP chemotherapy in	2040/27477/Long-term-outcome-of-patients-in-
DLBCL patients: A study by the Groupe d'Etudes des	<u>the-LNH-98-5</u>
Lymphomes de l'Adulte. <i>Blood</i> . 2010;116:2040-2045.	
Dunleavy K. Aggressive B cell lymphoma: Optimal	https://link.springer.com/article/10.1007/s11864-
therapy for MYC-positive, double-hit, and triple-hit	015-0374-0
DLBCL. Curr Treat Options Oncol. 2015;16:58.	
Inamdar A, et al. <i>Oncotarget.</i> 2016;7(30):48692-48731.	https://www.oncotarget.com/article/8961/text/
Lynch DT, et al. Mantle cell lymphoma. <i>StatPearls</i> . July 28, 2023.	https://www.ncbi.nlm.nih.gov/books/NBK536985/
National Cancer Institute (NCI). Cancer Stat Facts: NHL—	https://seer.cancer.gov/statfacts/html/follicular.ht
Follicular Lymphoma.	ml
Pavlasova G, et al. The regulation and function of CD20:	
an "enigma" of B-cell biology and targeted therapy.	https://haematologica.org/article/view/9425
Haematologica. 2020 Jun;105(6):1494-1506.	
Pescovitz MD. Rituximab, an anti-CD20 monoclonal	
antibody: History and mechanism of action. Am J	https://www.amjtransplant.org/article/S1600-
Transplant. 2006;6:859-866.	<u>6135(22)02663-6/fulltext</u>
Schmitz R, et al. Genetics and pathogenesis of diffuse	
large B-cell lymphoma. N Engl J Med. 2018;378:1396-	https://www.nejm.org/doi/10.1056/NEJMoa1801
1407.	<u>445</u>
1	

Bispecific Antibodies in B Cell NHL Therapy: Current State, Future Potential, and Adverse Event Profile

Reference	URL
AssayGenie. COVID-19 cytokine storm multiplex assays.	https://www.assaygenie.com/cytokine-storm-
	multiplex-assays
A Study Evaluating the Efficacy and Safety of Mosunetuzumab in Combination With Lenalidomide in Comparison to Rituximab in Combination With Lenalidomide in Patients With Follicular Lymphoma After at Least One Line of Systemic Therapy (Celestimo). ClinicalTrials.gov identifier: NCT04712097. Updated September 6, 2023.	https://classic.clinicaltrials.gov/ct2/show/NCT047120 97
A Study to Assess the Anti-Tumor Activity and Safety of Odronextamab in Patients With B-cell Non-Hodgkin Lymphoma That Have Been Previously Treated (ELM-2). ClinicalTrials.gov identifier: NCT03888105. Updated April 20, 2023.	https://classic.clinicaltrials.gov/ct2/show/NCT038881 05
Bannerji R, et al. Odronextamab (REGN1979), a human CD20 x CD3 bispecific antibody, induces durable, complete responses in patients with highly refractory B-cell non-Hodgkin lymphoma, including patients refractory to CAR T therapy. <i>Blood</i> . 2020;136(suppl 1):42-43.	https://www.sciencedirect.com/science/article/pii/SO 006497118696065
Bannerji R, et al. Odronextamab, a human CD20×CD3 bispecific antibody in patients with CD20-positive B-cell malignancies (ELM-1): Results from the relapsed or refractory non-Hodgkin lymphoma cohort in a single- arm, multicentre, phase 1 trial. <i>Lancet Haematol</i> . 2022;9:e327-e339.	https://www.thelancet.com/journals/lanhae/article/P IIS2352-3026(22)00072-2/fulltext
Budde LE, Assouline S, Sehn LH, et al. Single-agent mosunetuzumab shows durable complete responses in patients with relapsed or refractory B-cell lymphomas: Phase I dose-escalation study. <i>J Clin Oncol</i> . 2022;40:481-491.	https://ascopubs.org/doi/10.1200/JCO.21.00931
Budde LE, et al. Safety and efficacy of mosunetuzumab, a bispecific antibody, in patients with relapsed or refractory follicular lymphoma: A single-arm, multicentre, phase 2 study. <i>Lancet Oncol</i> . 2022;23:1055-1065.	<u>https://www.thelancet.com/journals/lanonc/article/P</u> <u>IIS1470-2045(22)00335-7/fulltext</u>
Chiu C, et al. Evaluation of pharmacodynamic biomarkers of epcoritamab (GEN3013; CD3XCD20): Results from a phase 1/2 dose-escalation study in patients with relapsed/refractory B-cell non-Hodgkin lymphoma. Presented at the 25 th Congress of the European Hematology Association; June 11-21, 2020. EHA Library. 2020:EP1330.	https://library.ehaweb.org/eha/2020/eha25th/29381 7/christopher.chiu.evaluation.of.pharmacodynamic.bi omarkers.of.epcoritamab.html

Reference	URL
Casulo C, et al. Early relapse of follicular lymphoma	
after rituximab plus cyclophosphamide, doxorubicin,	
vincristine, and prednisone defines patients at high risk	https://ascopubs.org/doi/10.1200/JCO.2014.59.7534
for death: An analysis from the National LymphoCare	
Study. J Clin Oncol. 2015;33:2516-2522.	
Dickinson MJ, et al. EHA 2020; Abstract S241	https://library.ehaweb.org/eha/2020/eha25th/29369
	0/michael.j.dickinson.cd20-
	tcb.in.relapsed.or.refractory.non-
	hodgkin.lymphoma.html
Dickinson MJ, et al. Glofitamab for relapsed or	
refractory diffuse large B-cell lymphoma. N Engl J Med.	https://www.nejm.org/doi/10.1056/NEJMoa2206913
2022;387:2220-2231.	
Dickinson M, et al. Glofitamab in patients with	
relapsed/refractory (R/R) diffuse large B-cell lymphoma	https://ascopubs.org/doi/abs/10.1200/JCO.2022.40.1
(DLBCL) and \geq 2 prior therapies: Pivotal phase II	6 suppl.7500
expansion results. J Clin Oncol. 2022;40(16 suppl):7500.	
Engelberts PJ, et al. DuoBody-CD3xCD20 induces potent	
T-cell-mediated killing of malignant B cells in preclinical	https://www.thelancet.com/journals/ebiom/article/P
models and provides opportunities for subcutaneous	IIS2352-3964(19)30840-0/fulltext
dosing. EBioMedicine. 2020;52:102625.	
González Barca E. Role of bispecific antibodies in	
relapsed/refractory diffuse large B-cell lymphoma in	https://www.frontiersin.org/articles/10.3389/fimmu.
the CART era. Front Immunol. 2022;13:909008.	<u>2022.909008/full</u>
Hosseini I, et al. Mitigating the risk of cytokine release	
syndrome in a Phase I trial of CD20/CD3 bispecific	https://www.asture.com/articles/a41540.020.00145
antibody mosunetuzumab in NHL: Impact of	https://www.nature.com/articles/s41540-020-00145-
translational system modeling. NPJ Syst Biol Appl.	<u>Z</u>
2020;6:28.	
Hutchings M, et al. Glofitamab, a novel, bivalent CD20-	
targeting T-cell-engaging bispecific antibody, induces	
durable complete remissions in relapsed or refractory	https://ascopubs.org/doi/10.1200/JCO.20.03175
B-cell lymphoma: A phase I trial. J Clin Oncol.	
2021;39:1959-1970.	
Hutchings M, et al. Subcutaneous epcoritamab induces	
complete responses with an encouraging safety profile	https://ashpublications.org/blood/article/136/Supple
across relapsed/refractory B-cell non-Hodgkin	ment%201/45/470017/Subcutaneous-Epcoritamab-
lymphoma subtypes, including patients with prior CAR-	Induces-Complete
T therapy: Updated dose escalation data. <i>Blood.</i>	
2020;136(suppl 1):45-46.	
Hutchings M, et al. Dose escalation of subcutaneous	https://www.tholopcot.com/journals/logget/article/D
epcoritamab in patients with relapsed or refractory B-	https://www.thelancet.com/journals/lancet/article/PI
cell non-Hodgkin lymphoma: An open-label, phase 1/2 study. <i>Lancet</i> . 2021;398:1157-1169.	IS0140-6736(21)00889-8/fulltext
fKim TM, et al. A phase 2 study of odronextamab	https://ashpublications.org/blood/article/136/Supple
(REGN1979), a CD20 x CD3 bispecific antibody, in patients with relapsed/refractory B-cell non-Hodgkin	ment%201/28/472688/A-Phase-2-Study-of-
lymphoma. <i>Blood</i> . 2020;136(suppl 1):28-29.	Odronextamab-REGN1979-a-CD20-x
iyinpiloina. Dood. 2020,130(suppi 1).20-23.	

Reference	URL
Lee DW, et al. Current concepts in the diagnosis and	https://ashpublications.org/blood/article/124/2/188/
management of cytokine release syndrome. <i>Blood</i> .	<u>32896/Current-concepts-in-the-diagnosis-and-</u>
2014;124:188-195.	management
	management
Lee DW, et al. ASTCT consensus grading for cytokine	
release syndrome and neurologic toxicity associated	https://www.astctjournal.org/article/S1083-
with immune effector cells. <i>Biol Blood Marrow</i>	8791(18)31691-4/fulltext
Transplant. 2019;25:625-638.	
Link BK, et al. Second-line and subsequent therapy and	
outcomes for follicular lymphoma in the United States:	https://onlinelibrary.wiley.com/doi/10.1111/bjh.1514
Data from the observational National LymphoCare	<u>9</u>
Study. Br J Haematol. 2019;184:660-663.	
Minson A, et al. Glofitamab CD20-TCB bispecific	https://www.tandfonline.com/doi/abs/10.1080/1042
antibody. Leuk Lymphoma. 2021;62:3098-3108.	<u>8194.2021.1953016</u>
National Comprehensive Cancer Network [®] . NCCN	https://www.poon.org/professionale/shusision_stated
Clinical Practice Guidelines in Oncology. B-Cell	https://www.nccn.org/professionals/physician_gls/pd
Lymphoma. Version 5.2023.	<u>f/b-cell.pdf</u>
Neelapu SS, et al. Chimeric antigen receptor T-cell	
therapy - assessment and management of toxicities.	https://www.nature.com/articles/nrclinonc.2017.148
Nat Rev Clin Oncol. 2018;15:47-62.	
Nie S, et al. Biology drives the discovery of bispecific	
antibodies as innovative therapeutics. <i>Antib Ther</i> .	https://academic.oup.com/abt/article/3/1/18/57392
2020;3:18-62.	<u>55</u>
Phillips T, et al. Glofitamab monotherapy induces high	
complete response rates in patients with heavily	https://medically.gene.com/global/en/unrestricted/h
pretreated relapsed or refractory mantle cell	aematology/ASH-2022/ash-2022-presentation-
lymphoma. Presented at the 64th ASH Annual Meeting;	phillips-glofitamab-monotherapy-i.html
December 10–13, 2022.	
Phillips T, et al. Glofitamab monotherapy induces high	https://ashpublications.org/blood/article/140/Supple
complete response rates in patients with heavily	ment%201/178/489039/Glofitamab-Monotherapy-
pretreated relapsed or refractory mantle cell	Induces-High-Complete
lymphoma. <i>Blood</i> . 2022;140(suppl 1):178-180.	
Schuster SJ, et al. Mosunetuzumab induces complete	
remissions in poor prognosis non-Hodgkin lymphoma	https://ashpublications.org/blood/article/134/Supple
patients, including those who are resistant to or	ment 1/6/427814/Mosunetuzumab-Induces-
relapsing after chimeric antigen receptor T-cell (CAR-T)	Complete-Remissions-in-Poor
therapies, and Is active in treatment through multiple	
lines. Blood. 2019;134(suppl 1):6.	
Smith EJ, et al. A novel, native-format bispecific	
antibody triggering T-cell killing of B-cells is robustly	https://www.nature.com/articles/srep17943
active in mouse tumor models and cynomolgus	
monkeys. <i>Sci Rep</i> . 2015;5:17943.	
Subklewe M. Blood Adv. 2021;5(2):607-612.	https://ashpublications.org/bloodadvances/article/5/
	2/607/475005/BiTEs-better-than-CAR-T-cells
Sun LL, et al. Anti-CD20/CD3 T cell-dependent bispecific	https://www.coionec.erg/dei/10/1126/a-it-set/a-it-
antibody for the treatment of B cell malignancies. Sci	https://www.science.org/doi/10.1126/scitransImed.a
Transl Med. 2015;7:287ra70.	<u>aa4802</u>

Reference	URL
Thieblemont C, et al. Epcoritamab, a novel, subcutaneous CD3xCD20 bispecific T-cell-engaging antibody, in relapsed or refractory large B-cell lymphoma: Dose expansion in a phase I/II Trial. <i>J Clin</i> <i>Oncol</i> . 2023;41:2238-2247.	https://ascopubs.org/doi/10.1200/JCO.22.01725
Wang M, et al. Acalabrutinib in relapsed or refractory mantle cell lymphoma (ACE-LY-004): A single-arm, multicentre, phase 2 trial. <i>Lancet</i> . 2018;391:659-667.	https://www.thelancet.com/journals/lancet/article/Pl IS0140-6736(17)33108-2/fulltext
Wang ML, et al. Long-term follow-up of MCL patients treated with single-agent ibrutinib: Updated safety and efficacy results. <i>Blood</i> . 2015;126:739-745.	https://ashpublications.org/blood/article/126/6/739/ 34612/Long-term-follow-up-of-MCL-patients-treated- with
Wang XV, et al. Measurable residual disease does not preclude prolonged progression-free survival in CLL treated with ibrutinib. <i>Blood</i> . 2021;138:2810-2827.	https://ashpublications.org/blood/article/138/26/281 0/476606/Measurable-residual-disease-does-not- preclude
Zhou K, et al. Zanubrutinib monotherapy in relapsed/refractory mantle cell lymphoma: A pooled analysis of two clinical trials. <i>J Hematol Oncol</i> . 2021;14:167.	https://jhoonline.biomedcentral.com/articles/10.118 6/s13045-021-01174-3
Zhu M, et al. Translational findings for odronextamab: From preclinical research to a first-in-human study in patients with CD20+ B-cell malignancies. <i>Clin Transl Sci</i> . 2022;15:954-966.	https://ascpt.onlinelibrary.wiley.com/doi/10.1111/cts .13212
Zugmaier G, et al. Long-term follow-up of serum immunoglobulin levels in blinatumomab-treated patients with minimal residual disease-positive B- precursor acute lymphoblastic leukemia. <i>Blood Cancer</i> <i>J</i> . 2014;4(9):244.	https://www.nature.com/articles/bcj201464