

Key Updates in Bispecific Antibodies in NHL Management—Unmet Needs and Opportunities for Improved Management in Later-Line Management

Toolkit

Overview of B-Cell Non-Hodgkin Lymphomas

Reference	URL
Abramson JS, Shipp MA. Advances in the biology and therapy of diffuse large B-cell lymphoma: Moving toward a molecularly targeted approach. <i>Blood</i> . 2005;106:1164-1174.	https://ashpublications.org/blood/article/106/4/1164/21561/Advances-in-the-biology-and-therapy-of-diffuse
Alizadeh AA, Eisen MB, Davis RE, et al. Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling. <i>Nature</i> . 2000;403:503-511.	https://www.nature.com/articles/35000501
American Cancer Society (ACS). Key statistics for non-Hodgkin lymphoma. Last revised January 17, 2024.	https://www.cancer.org/cancer/types/non-hodgkin-lymphoma/about/key-statistics.html
Armitage JO, Weisenburger DD. New approach to classifying non-Hodgkin's lymphomas: Clinical features of the major histologic subtypes. Non-Hodgkin's Lymphoma Classification Project. <i>J Clin Oncol</i> . 1998;16:2780-2795.	https://ascopubs.org/doi/10.1200/JCO.1998.16.8.2780
Chapuy B, Stewart C, Dunford AJ, et al. Molecular subtypes of diffuse large B cell lymphoma are associated with distinct pathogenic mechanisms and outcomes. <i>Nat Med</i> . 2018;24:1290-1291.	https://www.nature.com/articles/s41591-018-0016-8
Coiffier B, Lepage E, Brière J, et al. CHOP chemotherapy plus rituximab compared with CHOP alone in elderly patients with diffuse large-B-cell lymphoma. <i>N Engl J Med</i> . 2002;346:235-242.	https://www.nejm.org/doi/full/10.1056/NEJMoa011795
Coiffier B, Thieblemont C, Van Den Neste E, et al. Long-term outcome of patients in the LNH-98.5 trial, the first randomized study comparing rituximab-CHOP to standard CHOP chemotherapy in DLBCL patients: A study by the Groupe d'Etudes des Lymphomes de l'Adulte. <i>Blood</i> . 2010;116:2040-2045.	https://ashpublications.org/blood/article/116/12/2040/27477/Long-term-outcome-of-patients-in-the-LNH-98-5
de Leval L, Alizadeh AA, Bergsagel PL, et al. Genomic profiling for clinical decision making in lymphoid neoplasms. <i>Blood</i> . 2022;24;140:2193-2227.	https://ashpublications.org/blood/article/140/21/2193/486384/Genomic-profiling-for-clinical-decision-making-in
Dunleavy K. Aggressive B cell lymphoma: Optimal therapy for MYC-positive, double-hit, and triple-hit DLBCL. <i>Curr Treat Options Oncol</i> . 2015;16:58.	https://link.springer.com/article/10.1007/s11864-015-0374-0
Eyre TA, Savage KJ, Cheah CY, et al. CNS prophylaxis for diffuse large B-cell lymphoma. <i>Lancet Oncol</i> . 2022;23:e416-e426.	https://www.thelancet.com/journals/lanonc/article/PIIS1470-2045(22)00371-0/abstract
Leukemia and Lymphoma Society. NHL Subtypes.	https://www.lls.org/lymphoma/non-hodgkin-lymphoma/nhl-subtypes
Pescovitz MD. Rituximab, an anti-CD20 monoclonal antibody: History and mechanism of action. <i>Am J Transplant</i> . 2006;6:859-866.	https://www.amjtransplant.org/article/S1600-6135(22)02663-6/fulltext

Schmitz R, Wright GW, Huang DW, et al. Genetics and pathogenesis of diffuse large B-cell lymphoma. <i>N Engl J Med.</i> 2018;378:1396-1407.	https://www.nejm.org/doi/10.1056/NEJMoa1801445
Swerdlow SH, Campo E, Pileri SA, et al. The 2016 revision of the World Health Organization classification of lymphoid neoplasms. <i>Blood.</i> 2016;127:2375-2390.	https://ashpublications.org/blood/article/127/20/2375/35286/The-2016-revision-of-the-World-Health-Organization
International Non-Hodgkin's Lymphoma Prognostic Factors Project. A predictive model for aggressive non-Hodgkin's lymphoma. <i>N Engl J Med.</i> 1993;329:987-994.	https://www.nejm.org/doi/full/10.1056/nejm19939303291402
Ziepert M, Hasenclever D, Kuhnt E, et al. Standard International prognostic index remains a valid predictor of outcome for patients with aggressive CD20+ B-cell lymphoma in the rituximab era. <i>J Clin Oncol.</i> 2010;28:2373-2380.	https://ascopubs.org/doi/10.1200/JCO.2009.26.2493

Developments in Therapy for B-NHL

Reference	URL
Caimi PF, Ai WZ, Alderuccio JP, et al. Loncastuximab tesirine in relapsed/refractory diffuse large B-cell lymphoma: Long-term efficacy and safety from the phase 2 LOTIS-2 study. <i>Haematologica.</i> Published online August 31, 2023.	https://haematologica.org/article/view/haematol.2023.283459
Chiu C, Hiemstra IH, Johnson P, et al. Evaluation of pharmacodynamic biomarkers of epcoritamab (GEN3013; CD3XCD20): Results from a phase 1/2 dose-escalation study in patients with relapsed/refractory B-cell non-Hodgkin lymphoma. Presented at the 25 th Congress of the European Hematology Association; June 11-21, 2020. EHA Library. 2020:EP1330.	https://library.ehaweb.org/eha/2020/eha25th/293817/christopher.chiu.evaluation.of.pharmacodynamic.biomarkers.of.epcoritamab.html
Creative Biolabs. ADC Development Services for Non-Hodgkin Lymphoma (NHL) Research.	https://www.creative-biolabs.com/adc/adc-development-for-nhl.htm
Dickinson MJ, Carlo-Stella C, Morschhauser F, et al. Glofitamab for relapsed or refractory diffuse large B-cell lymphoma. <i>N Engl J Med.</i> 2022;387:2220-2231.	https://www.nejm.org/doi/10.1056/NEJMoa2206913
Dickinson M, Carlo-Stella C, Morschhauser F, et al. Glofitamab in patients with relapsed/refractory (R/R) diffuse large B-cell lymphoma (DLBCL) and ≥ 2 prior therapies: Pivotal phase II expansion results. <i>J Clin Oncol.</i> 2022;40(16 suppl):7500.	https://ascopubs.org/doi/abs/10.1200/JCO.2022.40.16_suppl.7500
Duell J, Maddocks KJ, González-Barca E, et al. Long-term outcomes from the phase II L-MIND study of tafasitamab (MOR208) plus lenalidomide in patients with relapsed or refractory diffuse large B-cell lymphoma. <i>Haematologica.</i> 2021;106:2417-2426.	https://haematologica.org/article/view/haematol.2020.275958
Engelberts PJ, Hiemstra IH, de Jong B, et al. DuoBody-CD3xCD20 induces potent T-cell-mediated killing of malignant B cells in preclinical models and provides opportunities for subcutaneous dosing. <i>EBioMedicine.</i> 2020;52:102625.	https://www.thelancet.com/journals/ebiom/article/PIIS2352-3964(19)30840-0/fulltext

González Barca E. Role of bispecific antibodies in relapsed/refractory diffuse large B-cell lymphoma in the CART era. <i>Front Immunol.</i> 2022;13:909008.	https://www.frontiersin.org/articles/10.3389/fimmu.2022.909008/full
Hutchings M, Morschhauser F, Iacoboni G, et al. Glofitamab, a novel, bivalent CD20-targeting T-cell-engaging bispecific antibody, induces durable complete remissions in relapsed or refractory B-cell lymphoma: A phase I trial. <i>J Clin Oncol.</i> 2021;39:1959-1970.	https://ascopubs.org/doi/10.1200/JCO.20.03175
Hutchings M, Mous R, Clausen MR, et al. Subcutaneous epcoritamab induces complete responses with an encouraging safety profile across relapsed/refractory B-cell non-Hodgkin lymphoma subtypes, including patients with prior CAR-T therapy: Updated dose escalation data. <i>Blood.</i> 2020;136(suppl 1):45-46.	https://ashpublications.org/blood/article/136/Supplement%201/45/470017/Subcutaneous-Epcoritamab-Induces-Complete
Hutchings M, Mous R, Clausen MR, et al. Dose escalation of subcutaneous epcoritamab in patients with relapsed or refractory B-cell non-Hodgkin lymphoma: An open-label, phase 1/2 study. <i>Lancet.</i> 2021;398:1157-1169.	https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(21)00889-8/fulltext
Kim TM, Alonso A, Prince M, et al. A phase 2 study of odronextamab (REGN1979), a CD20 x CD3 bispecific antibody, in patients with relapsed/refractory B-cell non-Hodgkin lymphoma. <i>Blood.</i> 2020;136(suppl 1):28-29.	https://ashpublications.org/blood/article/136/Supplement%201/28/472688/A-Phase-2-Study-of-Odronextamab-REGN1979-a-CD20-x
Lee DW, Gardner R, Porter DL, et al. Current concepts in the diagnosis and management of cytokine release syndrome. <i>Blood.</i> 2014;124:188-195.	https://ashpublications.org/blood/article/124/2/188/32896/Current-concepts-in-the-diagnosis-and-management
Lee DW, Santomasso BD, Locke FL, et al. ASTCT consensus grading for cytokine release syndrome and neurologic toxicity associated with immune effector cells. <i>Biol Blood Marrow Transplant.</i> 2019;25:625-638.	https://www.astctjournal.org/article/S1083-8791(18)31691-4/fulltext
Minson A, Dickinson M. Glofitamab CD20-TCB bispecific antibody. <i>Leuk Lymphoma.</i> 2021;62:3098-3108.	https://www.tandfonline.com/doi/abs/10.1080/10428194.2021.1953016
National Comprehensive Cancer Network®. NCCN Clinical Practice Guidelines in Oncology. B-Cell Lymphomas. Version 1.2024.	https://www.nccn.org/professionals/physician_gls/pdf/b-cell.pdf
Neelapu SS, Tummala S, Kebriaei P, et al. Chimeric antigen receptor T-cell therapy - assessment and management of toxicities. <i>Nat Rev Clin Oncol.</i> 2018;15:47-62.	https://www.nature.com/articles/nrclinonc.2017.148
Nie S, Wang Z, Moscoso-Castro M, et al. Biology drives the discovery of bispecific antibodies as innovative therapeutics. <i>Antib Ther.</i> 2020;3:18-62.	https://academic.oup.com/abt/article/3/1/18/5739255
Northend M, Wilson W, Osborne W, et al. Results of a United Kingdom real-world study of polatuzumab vedotin, bendamustine, and rituximab for relapsed/refractory DLBCL. <i>Blood Adv.</i> 2022;6:2920-2926.	https://ashpublications.org/bloodadvances/article/6/9/2920/483452/Results-of-a-United-Kingdom-real-world-study-of
Phillips T, Dickenson M, Morschhauser F, et al. Glofitamab monotherapy induces high complete response rates in patients with heavily pretreated relapsed or refractory mantle cell lymphoma. Presented at the 64th ASH Annual Meeting; December 10–13, 2022.	https://medically.gene.com/global/en/unrestricted/haematology/ASH-2022/ash-2022-presentation-phillips-glofitamab-monotherapy-i.html

Phillips T, Dickenson M, Morschhauser F, et al. Glofitamab monotherapy induces high complete response rates in patients with heavily pretreated relapsed or refractory mantle cell lymphoma. <i>Blood.</i> 2022;140(suppl 1):178-180.	https://ashpublications.org/blood/article/140/Supplement%201/178/489039/Glofitamab-Monotherapy-Induces-High-Complete
Qualls DA, Lambert N, Caimi PF, et al. Tafasitamab and lenalidomide in large B-cell lymphoma: Real-world outcomes in a multicenter retrospective study. <i>Blood.</i> 2023;142:2327-2331.	https://ashpublications.org/blood/article-abstract/142/26/2327/498041/Tafasitamab-and-lenalidomide-in-large-B-cell
Sehn LH, Herrera AF, Flowers CR, et al. Polatuzumab vedotin in relapsed or refractory diffuse large B-cell lymphoma. <i>J Clin Oncol.</i> 2020;38:155-165.	https://ascopubs.org/doi/10.1200/JCO.19.00172
Smith EJ, Olson K, Haber LJ, et al. A novel, native-format bispecific antibody triggering T-cell killing of B-cells is robustly active in mouse tumor models and cynomolgus monkeys. <i>Sci Rep.</i> 2015;5:17943.	https://www.nature.com/articles/srep17943
Sun LL, Ellerman D, Mathieu M, et al. Anti-CD20/CD3 T cell-dependent bispecific antibody for the treatment of B cell malignancies. <i>Sci Transl Med.</i> 2015;7:287ra70.	https://www.science.org/doi/10.1126/scitranslmed.aaa4802
Thieblemont C, Phillips T, Ghesquieres H, et al. Epcoritamab, a novel, subcutaneous CD3xCD20 bispecific T-cell-engaging antibody, in relapsed or refractory large B-cell lymphoma: Dose expansion in a phase I/II Trial. <i>J Clin Oncol.</i> 2023;41:2238-2247.	https://ascopubs.org/doi/10.1200/JCO.22.01725
Tilly H, Morschhauser F, Sehn LH, et al. Polatuzumab vedotin in previously untreated diffuse large B-cell lymphoma. <i>N Engl J Med.</i> 2022;386:351-363.	https://www.nejm.org/doi/full/10.1056/NEJMoa2115304

All URLs accessed March 14, 2024